

Mark Scheme (Results) January 2011

GCE

GCE Statistics S1 (6683) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2011

Publications Code UA026664

All the material in this publication is copyright

© Edexcel Ltd 2011

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M marks:** method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A marks:** Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B marks** are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol \surd will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- \square The second mark is dependent on gaining the first mark

January 2011
Statistics S1 6683
Mark Scheme

Question Number	Scheme	Marks
1.		
(a)	$S_{ll} = 327754.5 - \frac{4027^2}{50} = 3419.92$ $S_{lw} = 29330.5 - \frac{357.1 \times 4027}{50} = 569.666$	M1 A1 A1 (3)
(b)	$r = \frac{569.666}{\sqrt{3419.92 \times 289.6}} = 0.572$ awrt 0.572 or 0.573	M1 A1 (2)
(c)	As the length of the salmon increases the weight increases	B1ft (1) [6]
Notes		
(a)	M1 for at least one correct expression 1 st A1 for $S_{ll} =$ awrt 3420 (Condone $S_{xx} = \dots$ or even $S_{yy} = \dots$) 2 nd A1 for $S_{lw} =$ awrt 570 (Condone $S_{xy} = \dots$)	
(b)	M1 for attempt at correct formula. Must have their S_{ll} , S_{lw} and given S_{ww} in the correct places If S_{ll} , S_{lw} are correct and an answer of awrt 0.57 is seen then award M1A0 M0 for $\frac{29330.5}{\sqrt{327754.5 \times 289.6}}$	
(c)	B1ft for a comment mentioning “length” and “weight”, not just l and w , and the idea of longer salmon weighing more. e.g. “positive correlation between weight and length” is B0 since the idea of positive correlation is not explained. Allow “larger” instead of “heavier” or “longer” Ignore any spurious values mentioned such as 0.572 If their r is negative (but must be $r > -1$) ft an appropriate comment. Condone $r > 1$ if comment is correct. If $ r < 0.4$ allow a comment of no or little relationship between weight and length but for $0 < r < 0.4$ the printed answer is still acceptable too. Treat mention of “skewness” as ISW if a correct interpretation is given	

Question Number	Scheme	Marks
2.		
(a)	$2.8 + 5.6 + 2.3 + 9.4 + 0.5 + 1.8 + 84.6 = 107$ $\text{mean} = 107 / 28 (= 3.821\dots)$	M1 A1 (2)
(b)	It will have no effect since one is 4.5 under what it should be and the other is 4.5 above what it should be.	B1 dB1 (2) [4]
Notes		
(a)	M1 for a clear attempt to add the two sums. Accept a full expression or $2.8 + 5.6 + \dots + 84.6 = x$ where $100 < x < 110$ i.e. seeing at least two correct terms of Keith's and the 84.6 with a slip. A1 for awrt 3.8 (Condone 1 dp/2sf here since data is given to 1 dp or 2 sf) Accept $\frac{107}{28}$ or $3\frac{23}{28}$ or any exact equivalent Correct answer implies M1A1	
(b)	1 st B1 for clearly stating that it will have no effect. ("roughly the same" is B0 B0) 2 nd dB1 for a supporting reason that mentions the fact that the increase and decrease are the same and gives some numerical value(s) to support this. e.g. Sum of Keith's observations is still 22.4 (or mean is still 3.2) <u>or</u> Sum is still 107 <u>or</u> $9.4 - 4.9 = 5 - 0.5$ (o.e.) This second B1 is dependent on their saying there is no effect so B0B1 is not possible.	

Question Number	Scheme	Marks
3.	<p>(a) Outliers $14 + 1.5 \times (14 - 7) = 24.5$ $7 - 1.5 \times (14 - 7) = -3.5$</p> <p>Outlier 25 either upper limit acceptable on diagram</p>  <p style="text-align: right;">Sales in £'000</p>	<p>M1 A1</p> <p>M1 A1ft B1</p> <p style="text-align: right;">(5)</p>
(b)	Since $Q_3 - Q_2 < Q_2 - Q_1$. Allow written explanation negatively skew	<p>B1 dB1</p> <p style="text-align: right;">(2)</p>
(c)	not true since the lower quartile is 7000 and therefore 75% above 7000 not 10000 or 10 is inside the box or any other sensible comment	<p>B1 dB1</p> <p style="text-align: right;">(2) [9]</p>
Notes		
(a)	<p>A fully correct box-plot (either version) with no supporting work scores 5/5. Otherwise read on</p> <p>1st M1 for at least one correct calculation seen 1st A1 for 24.5 and -3.5 (or just negative noted) seen. May be read off the graph. If both values are seen but no calculation is given then M1A1, one value M1A0. 2nd M1 for a box with an upper and a lower whisker(s) with at least 2 correct values (condone no median marked) 2nd A1ft for 3, 7, 12, 14 and 20 or 24.5 in appropriate places and readable off their scale If both upper whiskers are seen A0 Apply ft for their <u>whiskers</u> being compatible with their <u>outlier limits</u> e.g. if their lower limit is + 3.5 then a lower whisker ending at 4 or 3.5 is OK B1 for only one outlier appropriately marked at 25</p> <p style="text-align: center;">Apply ± 0.5 square accuracy for diagram</p>	
(b)	<p>1st B1 for $Q_3 - Q_2 < Q_2 - Q_1$ statement or an equivalent statement in words Use of $Q_3 - Q_2 < Q_2 - Q_1$ does not require differences to be seen. 2nd dB1 for “negative skew” dependent on suitable reason given above. “correlation” is B0 “positive skew” with a supporting argument based on whiskers can score B1B1 e.g. “right hand whisker is longer than LH one so positive skew” $Q_3 - Q_2 < Q_2 - Q_1$ followed by “positive skew” is B1B0</p>	
(c)	<p>1st B1 for rejecting the company’s claim 2nd dB1 for an appropriate supporting reason. Dependent on rejecting company’s claim.</p>	

Question Number	Scheme	Marks
4.		
(a)	$b = \frac{1.688}{5.753} = 0.293$ $a = 3.22 - 4.42 \times 0.293 = 1.9231...$ $p = 1.92 + 0.293v$	M1A1 M1 A1 (4)
(b)	$v = \frac{85-5}{10} = 8$ $p = 1.92 + 0.293 \times 8 = 4.3$ <p style="text-align: right;">(awrt 4.3)</p>	M1 A1 (2) [6]
Notes		
(a)	<p style="text-align: center;">Can ignore (a) and (b) labels here</p> <p>1st M1 for a correct expression for b. $\frac{1.688}{1.168}$ is M0 1st A1 for awrt 0.29</p> <p>2nd M1 for use of $a = \bar{p} - b\bar{v}$ follow through their value of b (or even just the letter b) 2nd A1 for a complete equation with $a =$ awrt 1.92 and $b =$ awrt 0.293 y or $p = 1.92 + 0.293x$ is A0 Correct answer with no working is 4/4</p>	
(b)	<p>M1 for an attempt to find the value of v when $x = 85$ (at least 2 correct terms in $\pm \frac{85-5}{10}$) <u>or</u> for an attempt to find an equation for p in terms of x <u>and</u> using $x = 85$ Attempt at equation of p in x requires $p = 1.92 + 0.293 \frac{(x-5)}{10}$</p> <p>A1 for awrt 4.3 (award when first seen and apply ISW) N.B. $p = 1.92 + 0.293 \times 85$ (o.e.) is M0A0</p>	

Question Number	Scheme	Marks
5.		
(a)	Median = $32/2 = 16^{\text{th}}$ term (16.5) $\frac{x-39.5}{49.5-39.5} = \frac{16-14}{25-14}$ or $x = 39.5 + \left(\frac{2}{11} \times 10\right)$ Median = 41.3 (use of $n + 1$ gives 41.8) (awrt 41.3)	M1 A1 (2)
(b)	Mean = $\frac{1414}{32} = 44.1875$ (awrt 44.2) Standard deviation = $\sqrt{\frac{69378}{32} - \left(\frac{1414}{32}\right)^2}$ $= 14.7$ (or $s = 14.9$)	B1 M1 A1 (3)
(c)	mean > median therefore <u>positive skew</u>	B1ft B1ft (2) [7]
Notes		
(a)	M1 for an attempt to use interpolation to find the median. Condone use of 39 or 40 for 39.5 e.g. allow $39 + \frac{2}{11} \times 10$ (o.e.) or $40 + \frac{2}{11} \times 10$ (o.e.) to score M1A0 but must have the 10 A1 for awrt 41.3 (or awrt 41.8 if using $(n + 1)$)	
Mid-points	(b) B1 for awrt 44.2 M1 for a correct expression including square root. (Allow ft of their mean) A1 for awrt 14.7 (If using s for awrt 14.9) You may see $\sum t = 1339 \rightarrow \bar{t} = 41.8$ and $\sum t^2 = 62928 \rightarrow \sigma 14.7$ or $s = 14.9$ this scores B0 for the mean but can score M1 for a correct st.dev expression and A1 for ans. Correct answer only in (a) and (b) can score full marks but check $(n + 1)$ case in (a)	
Quartiles	(c) 1 st B1ft for a correct comparison of their mean and their median (may be in a formula) Calculating median – mean as negative is OK for this B1 but must say +ve skew for 2 nd B1 Only allow comparison to be ≈ 0 if $ \text{mean} - \text{median} \leq 0.5$ 2 nd B1ft for a correct description of skewness <u>based on their values of mean and median</u> . ft their values for mean and median not their previous calculation/comparison Must be compatible with their previous comparison (if they have one) “Positive skew” with no reason is B0B1 provided you can see their values that imply that. Description should be “positive” or “negative” or “no” skew or “symmetric” “Positive correlation” is B0 1 st B1ft if $Q_1 = \text{awrt } 32$ <u>and</u> $Q_3 = \text{awrt } 49$ seen and a correct comparison made. ft Q_2 2 nd B1ft if $Q_1 = \text{awrt } 32$ <u>or</u> $Q_3 = \text{awrt } 49$ seen and a correct description based on their quartiles and their comparison is made. (Should get “negative skew”)	

Question Number	Scheme									Marks															
6.																									
(a)	$k + 2k + 3k + 4k = 1$ or $10k = 1$ $k = 0.1$ (*) [allow verification with a comment e.g. “so $k = 0.1$ ”]									B1cso (1)															
(b)	$E(X) = 1 \times 0.1 + 2 \times 0.2 + 3 \times 0.3 + 4 \times 0.4 = 3$									M1 A1 (2)															
(c)	$E(X^2) = 1 \times 0.1 + 4 \times 0.2 + 9 \times 0.3 + 16 \times 0.4 = 10$									M1 A1 (2)															
(d)	$\text{Var}(X) = 10 - 9 (=1)$ $\text{Var}(2 - 5X) = 5^2 \text{Var}(X) = 25$									M1 M1 A1 (3)															
(e)	$P(1,3) + P(2,2) = 2 \times 0.1 \times 0.3 + 0.2 \times 0.2 = 0.1$ (*)									M1 A1cso (2)															
(f)	<table border="1"><tr><td>$X_1 + X_2$</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td></tr><tr><td>p</td><td>0.01</td><td>0.04</td><td>0.1</td><td>0.2</td><td>0.25</td><td>0.24</td><td>0.16</td></tr></table>	$X_1 + X_2$	2	3	4	5	6	7	8	p	0.01	0.04	0.1	0.2	0.25	0.24	0.16								B1 B1 (2)
$X_1 + X_2$	2	3	4	5	6	7	8																		
p	0.01	0.04	0.1	0.2	0.25	0.24	0.16																		
(g)	$P(2) + P(3) = 0.05$									M1A1 (2) [14]															

Question Number	Scheme	Marks
	Notes	
(a)	B1 for a clear attempt to use sum of probabilities = 1. Must see previous line as well as $k = 0.1$ A correct expression for $E(X)$ or $E(X^2)$ that is later divided by 4 scores M0	
(b)	M1 for a completely correct expression. May be implied by correct answer of 3 or $30k$ A1 for 3 only.	
(c)	M1 for a completely correct expression. May be implied by correct answer of 10 or $100k$ A1 for 10 only. [For $E(X^2) = 0.1 + 0.8 + 2.7 + 6.4 - 9 = 1$ scores M0A0 but accept this as $\text{Var}(X)$ in (d)]	
(d)	1 st M1 for using $\text{Var}(X) = E(X^2) - E(X)^2$, f.t. their values from (b) and (c) Allow this mark for $\text{Var}(X) = 10 - 9$ or better. May be implied if this is seen in (c). 2 nd M1 for $5^2 \text{Var}(X)$ or $25\text{Var}(X)$ can f.t. their $\text{Var}(X)$. Allow -5^2 if it later becomes $+25$ A1 for 25 only. Dependent upon both Ms Forming distribution for $Y = 2-5X$ gets M1 for $E(Y^2) = 194$ then M1A1 for $194 - 169 = 25$	
(e)	M1 for correctly identifying (1, 3) <u>or</u> (3, 1) <u>and</u> (2, 2) as required cases ($3k^2 + 4k^2$ or better) A1 cso for 0.1 only but must see evidence for M1	
(f)	1 st B1 for 0.2 correctly assigned. May be in table. 2 nd B1 for 0.16 correctly assigned. May be in table	
(g)	M1 for $P(2) + P(3)$. May be implied by correct answer of 0.05 A1 for 0.05 only. Correct answer only can score full marks in parts (b), (c), (f) and (g)	

Question Number	Scheme		Marks
7.	(a)		<p>both $\frac{2}{3}, \frac{1}{3}$ B1</p> <p>$\frac{4}{9}$ B1</p> <p>both $\frac{3}{5}, \frac{2}{5}$ B1</p> <p>all three of $\frac{4}{9}, \frac{4}{9}, \frac{5}{9}$ B1</p> <p>(4)</p>
(b)	$P(A) = P(RR) + P(YY) = \frac{1}{2} \times \frac{2}{5} + \frac{1}{2} \times \frac{2}{5} = \frac{2}{5}$	B1 for $\frac{1}{2} \times \frac{2}{5}$ (oe) seen at least once	B1 M1 A1 (3)
(c)	$P(B) = P(RRR) + P(RYR) + P(YRR) + P(YYR)$ $\left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3} \right) + \left(\frac{1}{2} \times \frac{3}{5} \times \frac{5}{9} \right) + \left(\frac{1}{2} \times \frac{3}{5} \times \frac{5}{9} \right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{4}{9} \right) = \frac{5}{9} (*)$	M1 for at least 1 case of 3 balls identified. (Implied by 2 nd M1)	M1
(d)	$P(A \cap B) = P(RRR) + P(YYR)$ $= \left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3} \right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{4}{9} \right) = \frac{2}{9} (*)$	M1 for identifying both cases and + probs. may be implied by correct expressions	M1
(e)	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$ $= \frac{2}{5} + \frac{5}{9} - \frac{2}{9} = \frac{11}{9}$	Must have some attempt to <u>use</u>	M1
			A1cao (2)

Question Number	Scheme		Marks
(f)	$\frac{P(RRR)}{P(RRR) + P(YYY)} = \frac{\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}}{\left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}\right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{5}{9}\right)} = \frac{6}{11}$	Probabilities must come from the product of 3 probs. from their tree diagram.	M1 A1ft A1 cao (3) [17]
	Notes		
(b)	M1 for both cases, and +, attempted, ft their values from tree diagram. May be 4 cases of 3 balls.		
(c)	2 nd M1 for all 4 correct expressions, ft their values from tree diagram. A1 is cso		
(e)	M1 for clear attempt to <u>use</u> the correct formula, must have some correct substitution. ft their (b)		
(f)	M1 for identifying the correct probabilities and forming appropriate fraction of probs. 1 st A1ft for a correct expression using probabilities from their tree Accept exact decimal equivalents. Correct answer only is full marks except in (c) and (d)		

Question Number	Scheme	Marks
8.		
(a)	$P(X > 168) = P\left(Z > \frac{168-160}{5}\right)$ $= P(Z > 1.6)$ $= 0.0548$ <p style="text-align: right;">awrt 0.0548</p>	M1 A1 A1 (3)
(b)	$P(X < w) = P\left(Z < \frac{w-160}{5}\right)$ $\frac{w-160}{5} = -2.3263$ $w = 148.37$ <p style="text-align: right;">awrt 148</p>	M1 B1 A1 (3)
(c)	$\frac{160-\mu}{\sigma} = 2.3263$ $\frac{152-\mu}{\sigma} = -1.2816$ $160-\mu = 2.3263\sigma$ $152-\mu = -1.2816\sigma$ $8 = 3.6079\sigma$ $\sigma = 2.21\dots$ $\mu = 154.84\dots$ <p style="text-align: right;">awrt 2.22 awrt 155</p>	M1 B1 B1 M1 A1 A1 (6) [12]
Notes		
(a)	M1 for an attempt to standardize 168 with 160 and 5 i.e. $\pm\left(\frac{168-160}{5}\right)$ or implied by 1.6 1 st A1 for $P(Z > 1.6)$ or $P(Z < -1.6)$ ie $z = 1.6$ and a correct inequality or 1.6 on a shaded diagram <p style="text-align: center;">Correct answer to (a) implies all 3 marks</p>	
(b)	M1 for attempting $\pm\left(\frac{w-160}{5}\right) = \text{recognizable } z \text{ value } (z > 1)$ B1 for $z = \pm 2.3263$ or better. Should be $z = \dots$ or implied so: $1 - 2.3263 = \frac{w-160}{5}$ is M0B0 A1 for awrt 148. This may be scored for other z values so M1B0A1 is possible <p style="text-align: center;">For awrt 148 only with no working seen award M1B0A1</p>	
(c)	M1 for attempting to standardize 160 or 152 with μ and σ (allow \pm) <u>and</u> equate to z value ($ z > 1$) 1 st B1 for awrt ± 2.33 or ± 2.32 seen 2 nd B1 for awrt ± 1.28 seen 2 nd M1 for attempt to solve their two linear equations in μ and σ leading to equation in just one variable 1 st A1 for $\sigma = \text{awrt } 2.22$. Award when 1 st seen 2 nd A1 for $\mu = \text{awrt } 155$. Correct answer only for part (c) can score all 6 marks. NB $\sigma = 2.21$ commonly comes from $z = 2.34$ and usually scores M1B0B1M1A0A1 <p style="text-align: center;">The A marks in (c) require both M marks to have been earned</p>	

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UA026664 January 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH