



- (a) the value of  $m$ ,

(4)

- (b) the value of  $a$ .

(4)

- (a) Find the value of  $V$ .

(b) find the acceleration of the car when its speed is  $20 \text{ m s}^{-1}$ .

This image shows a full page of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook paper. There are no margins, text, or other markings on the page.

- (c) the position vector of  $A$ .
- (5)**

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

A diagram showing a projectile launched from point A at a height  $h$  m above a horizontal ground line. The launch point A is marked with a vertical line segment of length  $h$  m down to the ground at point O. The projectile is launched at an angle  $\alpha^\circ$  to the horizontal with an initial speed of  $12 \text{ m s}^{-1}$ . The projectile follows a parabolic path and lands at point B on the ground.

The points  $O$  and  $B$  are on horizontal ground. The point  $A$  is  $h$  metres vertically above  $O$ . A particle  $P$  is projected from  $A$  with speed  $12 \text{ m s}^{-1}$  at an angle  $\alpha^\circ$  to the horizontal. The particle moves freely under gravity and hits the ground at  $B$ , as shown in Figure 1. The speed of  $P$  immediately before it hits the ground is  $15 \text{ m s}^{-1}$ .

- Given that 1.5 s after it is projected from A, P is at a point 4 m above the level of A, find

- (c) the direction of motion of  $P$  immediately before it reaches  $B$ .

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is no text or other markings on the paper.

The diagram shows an L-shaped figure with vertices labeled  $O$ ,  $A$ ,  $B$ ,  $C$ ,  $D$ , and  $E$ . The vertical segment  $OE$  has a length of  $4\text{ m}$ . The horizontal segment  $OA$  is divided into two parts:  $OC$  and  $CA$ . The length of  $OC$  is labeled as  $a\text{ m}$ , and the length of  $CA$  is labeled as  $4\text{ m}$ . The vertical segment  $CB$  has a length of  $a\text{ m}$ . A dashed line connects  $O$  to  $C$ .

### Figure 2


The uniform L-shaped lamina  $OABCDE$ , shown in Figure 2, is made from two identical rectangles. Each rectangle is 4 metres long and  $a$  metres wide. Giving each answer in terms of  $a$ , find the distance of the centre of mass of the lamina from

- $$\begin{aligned} \text{(a) } & OE, \\ \text{(b) } & OA. \end{aligned} \tag{4}$$

The lamina is freely suspended from  $O$  and hangs in equilibrium with  $OE$  at an angle  $\theta$  to the downward vertical through  $O$ , where  $\tan \theta = \frac{4}{3}$ .

- (c) Find the value of  $a$ . **(4)**

**6.**



### Figure 3

A uniform rod  $AB$  has weight  $30\text{ N}$  and length  $3\text{ m}$ . The rod rests in equilibrium on a rough horizontal peg  $P$  with its end  $A$  on smooth horizontal ground. The rod is in a vertical plane perpendicular to the peg. The rod is inclined at  $15^\circ$  to the ground and the point of contact between the peg and the rod is  $45\text{ cm}$  above the ground, as shown in Figure 3.

- (a) Show that the normal reaction at  $P$  has magnitude 25 N. (4)
- (b) Find the magnitude of the force on the rod at  $A$ . (4)

The coefficient of friction between the rod and the peg is  $\mu$ .

- (c) Find the range of possible values of  $\mu$ . (4)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Diagram showing two particles, P and Q, moving to the right. Particle P has mass  $m$  and velocity  $2u$ . Particle Q has mass  $2m$  and velocity  $3u$ .

### Figure 4

Two smooth particles  $P$  and  $Q$  have masses  $m$  and  $2m$  respectively. The particles are moving in the same direction in the same straight line, on a smooth horizontal plane, with  $Q$  in front of  $P$ . The particles are moving towards a fixed smooth vertical wall which is perpendicular to the direction of motion of the particles, as shown in Figure 4. The speed of  $P$  is  $2u$  and the speed of  $Q$  is  $3u$ . The coefficient of restitution between  $Q$  and the wall

is  $\frac{1}{3}$ . Particle  $Q$  strikes the wall, rebounds and then collides directly with  $P$ . The direction

of motion of each particle is reversed by this collision. Immediately after this collision the speed of  $P$  is  $v$  and the speed of  $Q$  is  $w$ .

(a) Show that  $v = 2w$ .

(5)

The total kinetic energy of  $P$  and  $Q$  immediately after they collide is half the total kinetic energy of  $P$  and  $Q$  immediately before they collide.

(b) Find the coefficient of restitution between  $P$  and  $Q$ .

(8)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.