

Version1.0



**General Certificate of Education (A-level)
January 2011**

Mathematics

MM2B

(Specification 6360)

Mechanics 2B

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Mark Scheme – General Certificate of Education (A-level) Mathematics – Mechanics 2B – January 2011

MM2B

Q	Solution	Marks	Total	Comments
1(a)	$\mathbf{r} = \int v \, dt$ $= (4t + t^3)\mathbf{i} + (12t - 4t^2)\mathbf{j} + \mathbf{c}$ <p>When $t = 0$, $\mathbf{r} = 5\mathbf{i} - 7\mathbf{j}$ $\mathbf{c} = 5\mathbf{i} - 7\mathbf{j}$</p> $\mathbf{r} = (5 + 4t + t^3)\mathbf{i} + (-7 + 12t - 4t^2)\mathbf{j}$	M1A1		M1 either \mathbf{i} or \mathbf{j} term correct. Condone no \mathbf{c}
		M1		Any attempt at \mathbf{c}
		A1	4	
(b)	$\mathbf{a} = \frac{dv}{dt}$ $\mathbf{a} = 6t\mathbf{i} - 8\mathbf{j}$	M1A1	2	M1 either term correct
(c)	Using $\mathbf{F} = m\mathbf{a}$ $\mathbf{F} = 2(6t\mathbf{i} - 8\mathbf{j})$ $= 12t\mathbf{i} - 16\mathbf{j}$ $\therefore \text{Magnitude of force is}$ $(144t^2 + 256)^{\frac{1}{2}} \text{ when } t = 1$ $= 20 \text{ N}$	M1 A1 M1 A1	4	Or: using $\mathbf{F} = m\mathbf{a}$ $\mathbf{F} = 2(6t\mathbf{i} - 8\mathbf{j})$ <p>When $t = 1$, $\mathbf{F} = 12\mathbf{i} - 16\mathbf{j}$</p> <p>Magnitude of force is $(144 + 256)^{\frac{1}{2}}$ $= 20 \text{ N}$</p>
Total			10	
2(a)	PE lost is $= 4 \times g \times 5 \cos 70$ $= 67.0 \text{ J}$	M1A1	2	M1 $4 \times g \times 5 \times \cos$ or $\sin 20$ or 70
(b)	KE is loss of PE \Rightarrow KE is 67.0 J	B1	1	ft
(c)	Using KE = $\frac{1}{2}mv^2$ $v^2 = 33.5$ <p>Speed of particle is 5.79 m s^{-1}</p>	M1 A1	2	(ft from (b))
Total			5	
3(a)	PE is $400 \times g \times 8$ $= 3200g \text{ [or } 31\,360]$	B1	1	
(b)	KE is $\frac{1}{2} \times 400 \times 2^2$ $= 800$	B1	1	
(c)	Work done per minute is 32 160 J <p>Power = $32\,160 \div 60$ $= 536 \text{ W}$</p>	M1 A1	2	$[(a) + (b)] \div 60$ CAO Accept 537 from 31 400 in (a)
Total			4	

Mark Scheme – General Certificate of Education (A-level) Mathematics – Mechanics 2B – January 2011

MM2B(cont)

Q	Solution	Marks	Total	Comments
4(a)	Moments about line AD : $5 \times 30 + 4 \times 10 = 9 \times \bar{x}$ $\bar{x} = \frac{190}{9}$ $= 21.1 \text{ cm}$	M1A1 A1	3	M1 2 of 3 terms correct
(b)	Moments about line AB : $5 \times 15 + 4 \times 25 = 9 \times \bar{y}$ $\bar{y} = \frac{175}{9}$ $\bar{y} = 19.4 \text{ cm}$	M1A1 A1	3	M1 2 of 3 terms correct If moments about DC ; 10.6 found SC2
(c)	$\tan \theta = \frac{80}{175}$ or $\frac{8.9}{19.4}$ $= 0.4571$ Angle is $\tan^{-1} 0.4571$ $= 24.6^\circ$	M1 A1 A1 A1	4	M1 use of tan A1 use of 8.9 or 80 ($30 - (a)$) Or 0.45876 $65.4^\circ \Rightarrow$ M1A1 only
(d)	Moments about the line PR : (or AD or BC) $30m = 4 \times 20$ or $9 \times \frac{80}{9}$ $m = \frac{8}{3}$	M1 A1 A1	3	
(e)	Centre of mass is at middle of lamina	E1	1	
	Total		14	

Mark Scheme – General Certificate of Education (A-level) Mathematics – Mechanics 2B – January 2011

MM2B(cont)

Q	Solution	Marks	Total	Comments
5(a)	Resolve vertically $R = mg$ If the particle is on the point of sliding, $F = \mu R$ $\therefore F = 0.3R = 0.3mg$ Resolving radially: $F = m\omega^2 r$ $0.3mg = m\omega^2 \times 0.8$ $\omega^2 = \frac{0.3 \times g}{0.8}$ $\omega = 1.92$	M1 A1 M1		Ignore all inequalities
(b)(i)	45 revolutions per minute = $\frac{90\pi}{60}$ $= \frac{3\pi}{2}$ or 4.71 radians per second	M1 A1	2	
(ii)	Resolving radially: $F = m\omega^2 r$ $m\mu g = m\left(\frac{3\pi}{2}\right)^2 \times 0.15$ $\mu = \frac{\left(\frac{3\pi}{2}\right)^2 \times 0.15}{g}$ $\mu = 0.340$	M1A1 A1		M1A1 either side correct A1 second side correct
	Total		4	CAO (accept 0.339)
			10	
6(a)	By conservation of energy $\frac{1}{2}m(5v)^2 = \frac{1}{2}m(3v)^2 + mg2a$ $8v^2 = 2ag$ $v = \sqrt{\frac{ag}{4}}$ or $\frac{1}{2}\sqrt{ag}$	M1 A1 A1		M1 for 3 terms , 2 KE and PE
(b)	Greatest and least values of tension are at the highest and lowest points of its path At top, $T = \frac{m(3v)^2}{a} - mg$ $= \frac{5}{4}mg$ At B, $T = \frac{m(5v)^2}{a} + mg$ $= \frac{29}{4}mg$ Ratio is 29 : 5	M1 A1ft M1 A1ft	4	ft - must be positive tension
	Total	A1	5	CAO Condone 5 : 29 or 1 : 5.8
			9	

MM2B(cont)

Q	Solution	Marks	Total	Comments
7(a)	Work done = $\int_0^e \frac{\lambda x}{l} dx$	M1	3	Condone lack of limits and 'dx'
	= $\left[\frac{\lambda x^2}{2l} \right]_0^e$	A1		Must include limits from integral
	= $\frac{\lambda e^2}{2l}$	A1		AG
(b)(i)	Using $T = \frac{\lambda x}{l}, 7g = \frac{196x}{2}$	M1	3	M1 could use 3g or 4g – at least 1 side correct
	$x = \frac{14g}{196}$	A1		
	= 0.7	A1		
(ii)	By C of Energy, when next at rest, EPE (initial) = PE change (for platform) + EPE (when at rest)		6	
	$\frac{196 \times 0.7^2}{2 \times 2} = 4 \times g \times (0.7 - x) + \frac{196x^2}{2 \times 2}$	M1A1 A1		M1 3 terms (not including $\frac{1}{2}mv^2$) A1 2 of 3 terms correct A1 all correct
	$2.45 = 2.8 - 4x + 5x^2$	m1		
	$100x^2 - 80x + 7 = 0$	A1		
	$(10x - 7)(10x - 1) = 0$ $x = 0.1$	A1		[last A1, must give 0.1, not 0.1 and 0.7]
(b)(ii)	Alternative			(where X is distance moved upwards)
	$\frac{196 \times 0.7^2}{2 \times 2} = 4gX + \frac{196(0.7 - X)^2}{2 \times 2}$	(M1) (A1) (A1)		
	$4gX = 98 \times 0.7X + 49X^2$	(m1)		
	$X = 0, 0.6$	(A1A1)		
(iii)	Max speed when $T = mg$	M1	3	Or mid-point of values 0.2 and 0.6 above SC2
	$4g = \frac{196x}{2}$	A1		
	$x = 0.4$	A1		
Total			15	

Mark Scheme – General Certificate of Education (A-level) Mathematics – Mechanics 2B – January 2011

MM2B

Q	Solution	Marks	Total	Comments
8(a)(i)	$F = 65g - 260v$ $= 65(9.8 - 4v)$	B1	1	Accept $260v - 65g$ AG must see $65g$ or 260
(ii)	Using $F = ma$ $65 \frac{dv}{dt} = 65(9.8 - 4v)$ $\frac{dv}{dt} = -4(v - 2.45)$	M1 A1	2	Need to see terms in m (condone $-$ sign) AG
(b)	$\frac{1}{v - 2.45} \frac{dv}{dt} = -4$ $\int \frac{1}{v - 2.45} dv = -\int 4 dt$ $\ln(v - 2.45) = -4t + c$ $v - 2.45 = Ce^{-4t}$ $t = 0, v = 19.6$ $\therefore C = 17.15$ or $e^{2.84}$ $\therefore v = 2.45 + 17.15e^{-4t}$ $2.45 + 17.2e^{-4t}$	B1 M1 A1 A1 A1	5	M1 log side correct $-4t + c$ Or $c = \ln 17.15$ or 2.84
	Total		8	
	TOTAL		75	