

1	Direction of $l_1 = k[7, 0, -10]$ } Direction of $l_2 = k[1, 3, -1]$ }	B1	For both directions
	<i>EITHER</i> $\mathbf{n} = [7, 0, -10] \times [1, 3, -1]$	M1	For finding vector product of directions of l_1 and l_2
	<i>OR</i> $\begin{cases} [x, y, z] \cdot [7, 0, -10] = 0 \Rightarrow 7x - 10z = 0 \\ [x, y, z] \cdot [1, 3, -1] = 0 \Rightarrow x + 3y - z = 0 \end{cases}$ $\Rightarrow \mathbf{n} = k[10, -1, 7]$	A1	<i>OR</i> for using 2 scalar products and obtaining equations For correct \mathbf{n}
METHOD 1			
Vector $(\mathbf{a} - \mathbf{b})$ from l_1 to $l_2 = \pm[4, 6, -10]$	B1	For a correct vector	
<i>OR</i> $\pm[-4, 3, 1]$ <i>OR</i> $\pm[3, 3, -9]$ <i>OR</i> $\pm[-3, 6, 0]$	M1*	For finding $(\mathbf{a} - \mathbf{b}) \cdot \mathbf{n}$	
$d = \frac{ (\mathbf{a} - \mathbf{b}) \cdot \mathbf{n} }{ \mathbf{n} } = \frac{36}{\sqrt{150}}$	M1 (*dep)	For $ \mathbf{n} $ in denominator <i>OR</i> for using $\hat{\mathbf{n}}$	
$d = \frac{6}{5}\sqrt{6} \approx 2.94$	A1	For correct distance AEF	
METHOD 2 Planes containing l_1 and l_2 perp. to \mathbf{n}			
are $\mathbf{r} \cdot [10, -1, 7] = p_1 = 70$, $\mathbf{r} \cdot [10, -1, 7] = p_2 = 34$	M1*	For finding planes and $p_1 - p_2$ seen	
$\Rightarrow d = \frac{ 70 - 34 }{\sqrt{150}} = \frac{36}{\sqrt{150}} = \frac{6}{5}\sqrt{6} \approx 2.94$	B1	For $p_1 = 70k$ and $p_2 = 34k$	
	M1 (*dep)	For $ \mathbf{n} $ in denominator <i>OR</i> for using $\hat{\mathbf{n}}$	
	A1	For correct distance AEF	
METHOD 3			
$\mathbf{r}_1 = [7\lambda, 0, 10 - 10\lambda]$ <i>OR</i> $[7 + 7\lambda, 0, -10\lambda]$	B1	For correct points on l_1 and l_2	
$\mathbf{r}_2 = [4 + \mu, 6 + 3\mu, -\mu]$ <i>OR</i> $[3 + \mu, 3 + 3\mu, 1 - \mu]$		using different parameters	
$\begin{array}{l} 7\lambda + 10\alpha - \mu = \begin{vmatrix} 4 & -3 & 3 & -4 \\ -\alpha - 3\mu & 6 & 3 & 3 \\ -10\lambda + 7\alpha + \mu & -10 & 0 & -9 & 1 \end{vmatrix} \\ \Rightarrow \alpha = -\frac{6}{25} \end{array}$	M1*	For setting up 3 linear equations from $\mathbf{r}_1 + \alpha\mathbf{n} = \mathbf{r}_2$ and solving for α	
$ \mathbf{n} = \sqrt{150}$	M1 (*dep)	For $ \mathbf{n} $ seen multiplying α	
$\Rightarrow d = \frac{6}{25}\sqrt{150} = \frac{6}{5}\sqrt{6} \approx 2.94$	A1	For correct distance AEF	
7			

2 (i)	$ar = r^5a \Rightarrow rar = r^6a$ $r^6 = e \Rightarrow rar = a$	M1 A1	Pre-multiply $ar = r^5a$ by r 2 Use $r^6 = e$ and obtain answer AG

(ii)	METHOD 1 For $n = 1, rar = a$ OR For $n = 0, r^0 ar^0 = a$ Assume $r^k ar^k = a$ EITHER Assumption $\Rightarrow r^{k+1} ar^{k+1} = rar = a$ OR $r^{k+1} ar^{k+1} = r.r^k ar^k.r = rar = a$ OR $r^{k+1} ar^{k+1} = r^k.rar.r^k = r^k ar^k = a$ Hence true for all $n \in \mathbb{Z}^+$	B1 M1 A1 A1	For stating true for $n = 1$ OR for $n = 0$ For attempt to prove true for $k + 1$ For obtaining correct form 4 For statement of induction conclusion

	METHOD 2 $r^2 ar^2 = r.rar.r = rar = a$, similarly for $r^3 ar^3 = a$ $r^4 ar^4 = r.r^3 ar^3.r = rar = a$, similarly for $r^5 ar^5 = a$ $r^6 ar^6 = eae = a$ For $n > 6, r^n = r^{n \bmod 6}$, hence true for all $n \in \mathbb{Z}^+$	M1 A1 B1 A1	For attempt to prove for $n = 2, 3$ For proving true for $n = 2, 3, 4, 5$ For showing true for $n = 6$ For using $n \bmod 6$ and correct conclusion

	METHOD 3 $r^n ar^n = r^{n-1}.rar.r^{n-1}$ OR $r^n ar^n = r^n.r^5 ar^{n-1} = r^{n+5} ar^{n-1}$ $= r^{n-1} ar^{n-1}$ $= r^{n-2} ar^{n-2} = \dots$ $= rar = a$	M1 A1 A1 B1	Starting from n , for attempt to prove true for $n - 1$ For proving true for $n - 1$ For continuation from $n - 2$ downwards For final use of $rar = a$ SR can be done in reverse

	METHOD 4 $ar = r^5a \Rightarrow ar^2 = r^5 ar = r^{10}a$ etc. $\Rightarrow ar^n = r^{5n}a$ $\Rightarrow r^n ar^n = r^{6n}a$ $= ea = a$	M1 A1 B1 A1	For attempt to derive $ar^n = r^{5n}a$ For correct equation SR may be stated without proof For pre-multiplication by r^n For obtaining a ($r^6 = e$ may be implied)


6			

3

(i) $w^2 = \cos \frac{4}{5}\pi + i \sin \frac{4}{5}\pi$
 $w^3 = \cos \frac{6}{5}\pi + i \sin \frac{6}{5}\pi$
 $w^* = \cos \frac{2}{5}\pi - i \sin \frac{2}{5}\pi$
 $= \cos \frac{8}{5}\pi + i \sin \frac{8}{5}\pi$

Allow $\text{cis } \frac{k}{5}\pi$ and $e^{\frac{k}{5}\pi i}$ throughout
 B1 For correct value
 B1 For correct value
 B1 For w^* seen or implied
 B1 4 For correct value
SR For exponential form with i missing, award B0 first time, allow others

(ii)



B1* For $1+w$ in approximately correct position
 B1 For $AB \approx BC \approx CD$
 (*dep)
 B1 For BC, CD equally inclined to Im axis
 (*dep)
 B1 4 For E at the origin
 Allow points joined by arcs, or not joined
 Labels not essential

(iii) $z^5 - 1 = 0$ OR $z^5 + z^4 + z^3 + z^2 + z = 0$

B1 1 For correct equation **AEF** (in any variable)
 Allow factorised forms using w , exp or trig

9

4 (i)

$y = xz \Rightarrow \frac{dy}{dx} = z + x \frac{dz}{dx}$
 $\Rightarrow xz + x^2 \frac{dz}{dx} - xz = x \cos z \Rightarrow x \frac{dz}{dx} = \cos z$
 $\Rightarrow \int \sec z \, dz = \int \frac{1}{x} \, dx$
 $\Rightarrow \ln(\sec z + \tan z) = \ln kx$
 OR $\ln \tan\left(\frac{1}{2}z + \frac{1}{4}\pi\right) = \ln kx$
 $\Rightarrow \sec\left(\frac{y}{x}\right) + \tan\left(\frac{y}{x}\right) = kx$
 OR $\tan\left(\frac{y}{2x} + \frac{1}{4}\pi\right) = kx$

B1 For correct differentiation of substitution
 M1 For substituting into DE
 A1 For DE in variables separable form
 M1 For attempt at integration to \ln form on LHS
 A1 For correct integration (k not required here)
 A1 6 For correct solution
AEF including $\text{RHS} = e^{(\ln x)+c}$

(ii) $(4, \pi) \Rightarrow \sec \frac{1}{4}\pi + \tan \frac{1}{4}\pi = 4k$

OR $\tan\left(\frac{1}{8}\pi + \frac{1}{4}\pi\right) = 4k$

$\Rightarrow \sec\left(\frac{y}{x}\right) + \tan\left(\frac{y}{x}\right) = \frac{1}{4}(1 + \sqrt{2})x$

OR $\tan\left(\frac{y}{2x} + \frac{1}{4}\pi\right) = \left(\frac{1}{4} \tan \frac{3}{8}\pi\right)x$ or $\frac{1}{4}(1 + \sqrt{2})x$

M1 For substituting $(4, \pi)$ into their solution (with k)
 A1 2 For correct solution **AEF**
 Allow decimal equivalent 0.60355 x
 Allow $e^{\ln x}$ for x

8

<p>5 (i) $C + iS = 1 + \frac{1}{2}e^{i\theta} + \frac{1}{4}e^{2i\theta} + \frac{1}{8}e^{3i\theta} + \dots$</p>	<p>M1</p>	<p>For using $\cos n\theta + i \sin n\theta = e^{in\theta}$ at least once for $n \geq 2$</p>
$= \frac{1}{1 - \frac{1}{2}e^{i\theta}} = \frac{2}{2 - e^{i\theta}}$	<p>A1</p>	<p>For correct series</p>
	<p>M1</p>	<p>For using sum of infinite GP</p>
	<p>A1</p>	<p>4 For correct expression AG SR For omission of 1st stage award up to M0 A0 M1 A1 OEW</p>

<p>(ii) $C + iS = \frac{2(2 - e^{-i\theta})}{(2 - e^{i\theta})(2 - e^{-i\theta})}$</p>	<p>M1</p>	<p>For multiplying top and bottom by complex conjugate</p>
$= \frac{4 - 2e^{-i\theta}}{4 - 2(e^{i\theta} + e^{-i\theta}) + 1} = \frac{4 - 2\cos\theta + 2i\sin\theta}{4 - 4\cos\theta + 1}$	<p>M1</p>	<p>For reverting to $\cos\theta$ and $\sin\theta$ and equating Re OR Im parts</p>
$\Rightarrow C = \frac{4 - 2\cos\theta}{5 - 4\cos\theta}, S = \frac{2\sin\theta}{5 - 4\cos\theta}$	<p>A1</p>	<p>For correct expression for C AG</p>
	<p>A1</p>	<p>4 For correct expression for S</p>

8		
<p>6 (i) Aux. equation $m^2 + 2m + 17 = 0$</p>	<p>M1</p>	<p>For attempting to solve correct auxiliary equation</p>
$\Rightarrow m = -1 \pm 4i$	<p>A1</p>	<p>For correct roots</p>
<p>CF $(y =) e^{-x} (A \cos 4x + B \sin 4x)$</p>	<p>A1√</p>	<p>For correct CF (allow $A \frac{\cos}{\sin} (4x + \varepsilon)$) (trig terms required, not $e^{\pm 4ix}$) f.t. from their m with 2 arbitrary constants</p>
<p>PI $(y =) px + q \Rightarrow 2p + 17(px + q) = 17x + 36$</p>	<p>M1</p>	<p>For stating and substituting PI of correct form</p>
$\Rightarrow p = 1$	<p>A1</p>	<p>For correct value of p</p>
<p>and $q = 2$</p>	<p>A1</p>	<p>For correct value of q</p>
<p>GS $y = e^{-x} (A \cos 4x + B \sin 4x) + x + 2$</p>	<p>B1√</p>	<p>7 For GS. f.t. from their CF+PI with 2 arbitrary constants in CF and none in PI. Requires $y =$.</p>

<p>(ii) $x \gg 0 \Rightarrow e^{-x} \rightarrow 0$ OR very small</p>	<p>B1</p>	<p>For correct statement. Allow graph</p>
$\Rightarrow y = x + 2$ approximately	<p>B1√</p>	<p>2 For correct equation Allow \approx, \rightarrow and in words Allow relevant f.t. from linear part of GS</p>

9		

7 (i)	$(1, 3, 5)$ and $(5, 2, 5) \Rightarrow \pm[4, -1, 0]$ in Π	M1	For finding a vector in Π
	$\mathbf{n} = [2, -2, 3] \times [4, -1, 0] = k[1, 4, 2]$	M1	For finding vector product of direction vectors of l and a line in Π
	$\Rightarrow \mathbf{r} \cdot [1, 4, 2] = 23$	A1	For correct \mathbf{n}
		A1	4 For correct equation. Allow multiples
(ii)	METHOD 1		
	Perpendicular to Π through $(-7, -3, 0)$ meets Π	M1	For using perpendicular from point on l to Π Award mark for $k\mathbf{n}$ used
	where $(-7+k)+4(-3+4k)+2(2k)=23$	M1	For substituting parametric line coords into Π
	$\Rightarrow k=2 \Rightarrow d=2\sqrt{1^2+4^2+2^2}=2\sqrt{21} \approx 9.165$	M1	For normalising the \mathbf{n} used in this part
		A1	4 For correct distance AEF
	METHOD 2		
	Π is $x+4y+2z=23$	M1	For attempt to use formula for perpendicular distance
	$\Rightarrow d = \frac{ (-7)+4(-3)+2(0)-23 }{\sqrt{1^2+4^2+2^2}} = 2\sqrt{21} \approx 9.165$	M1	For substituting a point on l into plane equation
		M1	For normalising the \mathbf{n} used in this part
		A1	For correct distance AEF
METHOD 3			
$\mathbf{m} = [1, 3, 5] - [-7, -3, 0] = (\pm)[8, 6, 5]$	M1	For finding a vector from l to Π	
$OR = [5, 2, 5] - [-7, -3, 0] = (\pm)[12, 5, 5]$			
$\Rightarrow d = \frac{\mathbf{m} \cdot [1, 4, 2]}{\sqrt{1^2+4^2+2^2}} = \frac{42}{\sqrt{21}} = 2\sqrt{21} \approx 9.165$	M1	For finding $\mathbf{m} \cdot \mathbf{n}$	
	M1	For normalising the \mathbf{n} used in this part	
	A1	For correct distance AEF	
METHOD 4			
$[-7, -3, 0] + k[1, 4, 2] = [1, 3, 5] + s[2, -2, 3] + t[4, -1, 0]$	M1	As Method 1, using parametric form of Π For using perpendicular from point on l to Π Award mark for $k\mathbf{n}$ used	
$\left. \begin{matrix} k-2s-4t=8 \\ 4k+2s+t=6 \\ 2k-3s=5 \end{matrix} \right\} \Rightarrow k=2 \left(s=-\frac{1}{3}, t=-\frac{4}{3} \right)$	M1	For setting up and solving 3 equations	
$\Rightarrow d = 2\sqrt{1^2+4^2+2^2} = 2\sqrt{21} \approx 9.165$	M1	For normalising the \mathbf{n} used in this part	
	A1	For correct distance AEF	
METHOD 5			
$d_1 = \frac{23}{\sqrt{1^2+4^2+2^2}} = \frac{23}{\sqrt{21}}$	M1	For attempt to find distance from O to Π OR from O to parallel plane containing l	
$d_2 = \frac{[-7, -3, 0] \cdot [1, 4, 2]}{\sqrt{1^2+4^2+2^2}} = \frac{-19}{\sqrt{21}}$	M1	For normalising the \mathbf{n} used in this part	
$\Rightarrow d_1 - d_2 = d = \frac{23 - (-19)}{\sqrt{21}} = 2\sqrt{21} \approx 9.165$	M1	For finding $d_1 - d_2$	
	A1	For correct distance AEF	
(iii)	$(-7, -3, 0) + k(1, 4, 2)$	M1	State or imply coordinates of a point on the reflected line
	Use $k=4$	M1	State or imply $2 \times$ distance from (ii) Allow $k = \pm 4$ OR $\pm 4\sqrt{21}$ f.t. from (ii)
	$\mathbf{b} = [2, -2, 3]$	B1	For stating correct direction
	$\mathbf{a} = [-3, 13, 8]$	A1	4 For correct point seen in equation $\mathbf{r} = \mathbf{a} + t\mathbf{b}$
	$\mathbf{r} = [-3, 13, 8] + t[2, -2, 3]$	A1	AEF in this form

8 (i)	$\{A, D\}$ OR $\{A, E\}$ OR $\{A, F\}$	B1	1	For stating any one subgroup																																																	
(ii)	A is the identity 5 is not a factor of 6 OR elements can be only of order 1, 2, 3, 6	B1 B1	2	For identifying A as the identity For reference to factors of 6																																																	
(iii)	$BE = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = D$, $EB = \begin{pmatrix} 0 & \omega \\ \omega^2 & 0 \end{pmatrix} = F$ D or $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, F or $\begin{pmatrix} 0 & \omega \\ \omega^2 & 0 \end{pmatrix} \in M$ \Rightarrow closure property satisfied	M1 A1 A1 A1	4	For finding BE and EB AND using $\omega^3 = 1$ For correct BE (D or matrix) For correct EB (F or matrix) For justifying closure																																																	
(iv)	$B^{-1} = \frac{1}{1} \begin{pmatrix} \omega^2 & 0 \\ 0 & \omega \end{pmatrix} = C$ $E^{-1} = \frac{1}{-1} \begin{pmatrix} 0 & -\omega^2 \\ -\omega & 0 \end{pmatrix} = E$	M1 A1 A1	3	For correct method of finding either inverse For correct $B^{-1} = C$ Allow $\begin{pmatrix} \omega^2 & 0 \\ 0 & \omega \end{pmatrix}$ For correct $E^{-1} = E$ Allow $\begin{pmatrix} 0 & \omega^2 \\ \omega & 0 \end{pmatrix}$																																																	
(v)	METHOD 1 M is not commutative e.g. from $BE \neq EB$ in part (iii) N is commutative (as $\times \pmod 9$ is commutative) \Rightarrow M and N not isomorphic	B1 B1 B1#	3	For justification of M being not commutative For statement that N is commutative For correct conclusion																																																	
	METHOD 2 Elements of M have orders 1, 3, 3, 2, 2, 2 Elements of N have orders 1, 6, 3, 2, 3, 6 Different orders OR self-inverse elements \Rightarrow M and N not isomorphic	B1* B1 (*dep) B1#		For all orders of one group correct For sufficient orders of the other group correct For correct conclusion SR Award up to B1 B1 B1 if the self-inverse elements are sufficiently well identified for the groups to be non-isomorphic																																																	
	METHOD 3 M has no generator since there is no element of order 6 N has 2 OR 5 as a generator \Rightarrow M and N not isomorphic	B1 B1 B1#		For all orders of M shown correctly For stating that N has generator 2 OR 5 For correct conclusion																																																	
	METHOD 4 <table border="1" style="display: inline-table; vertical-align: middle;"> <thead> <tr> <th>M</th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> <th>E</th> <th>F</th> </tr> </thead> <tbody> <tr> <td>A</td> <td>A</td> <td>B</td> <td>C</td> <td>D</td> <td>E</td> <td>F</td> </tr> <tr> <td>B</td> <td>B</td> <td>C</td> <td>A</td> <td>F</td> <td>D</td> <td>E</td> </tr> <tr> <td>C</td> <td>C</td> <td>A</td> <td>B</td> <td>E</td> <td>F</td> <td>D</td> </tr> <tr> <td>D</td> <td>D</td> <td>E</td> <td>F</td> <td>A</td> <td>B</td> <td>C</td> </tr> <tr> <td>E</td> <td>E</td> <td>F</td> <td>D</td> <td>C</td> <td>A</td> <td>B</td> </tr> <tr> <td>F</td> <td>F</td> <td>D</td> <td>E</td> <td>B</td> <td>C</td> <td>A</td> </tr> </tbody> </table> N 1 2 4 8 7 5 1 1 2 4 8 7 5 2 2 4 8 7 5 1 4 4 8 7 5 1 2 8 8 7 5 1 2 4 7 7 5 1 2 4 8 5 5 1 2 4 8 7 \Rightarrow M and N not isomorphic	M	A	B	C	D	E	F	A	A	B	C	D	E	F	B	B	C	A	F	D	E	C	C	A	B	E	F	D	D	D	E	F	A	B	C	E	E	F	D	C	A	B	F	F	D	E	B	C	A	B1* B1 (*dep) B1#		For stating correctly all 6 squared elements of one group For stating correctly sufficient squared elements of the other group For correct conclusion
M	A	B	C	D	E	F																																															
A	A	B	C	D	E	F																																															
B	B	C	A	F	D	E																																															
C	C	A	B	E	F	D																																															
D	D	E	F	A	B	C																																															
E	E	F	D	C	A	B																																															
F	F	D	E	B	C	A																																															
				# In all Methods, the last B1 is dependent on at least one preceding B1																																																	