

General Certificate of Education
January 2008
Advanced Level Examination



MATHEMATICS
Unit Further Pure 2

MFP2

Thursday 31 January 2008 9.00 am to 10.30 am

For this paper you must have:

- an 8-page answer book
 - the blue AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MFP2.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

Information

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

Answer **all** questions.

1 (a) Express $4 + 4i$ in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. (3 marks)

(b) Solve the equation

$$z^5 = 4 + 4i$$

giving your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. (5 marks)

2 (a) Show that

$$(2r + 1)^3 - (2r - 1)^3 = 24r^2 + 2 \quad (3 \text{ marks})$$

(b) Hence, using the method of differences, show that

$$\sum_{r=1}^n r^2 = \frac{1}{6}n(n+1)(2n+1) \quad (6 \text{ marks})$$

3 A circle C and a half-line L have equations

$$|z - 2\sqrt{3} - i| = 4$$

and
$$\arg(z + i) = \frac{\pi}{6}$$

respectively.

(a) Show that:

(i) the circle C passes through the point where $z = -i$; (2 marks)

(ii) the half-line L passes through the centre of C . (3 marks)

(b) On one Argand diagram, sketch C and L . (4 marks)

(c) Shade on your sketch the set of points satisfying both

$$|z - 2\sqrt{3} - i| \leq 4$$

and
$$0 \leq \arg(z + i) \leq \frac{\pi}{6} \quad (2 \text{ marks})$$

4 The cubic equation

$$z^3 + iz^2 + 3z - (1 + i) = 0$$

has roots α , β and γ .

(a) Write down the value of:

(i) $\alpha + \beta + \gamma$; *(1 mark)*

(ii) $\alpha\beta + \beta\gamma + \gamma\alpha$; *(1 mark)*

(iii) $\alpha\beta\gamma$. *(1 mark)*

(b) Find the value of:

(i) $\alpha^2 + \beta^2 + \gamma^2$; *(3 marks)*

(ii) $\alpha^2\beta^2 + \beta^2\gamma^2 + \gamma^2\alpha^2$; *(4 marks)*

(iii) $\alpha^2\beta^2\gamma^2$. *(2 marks)*

(c) Hence write down a cubic equation whose roots are α^2 , β^2 and γ^2 . *(2 marks)*

5 Prove by induction that for all integers $n \geq 1$

$$\sum_{r=1}^n (r^2 + 1)(r!) = n(n + 1)! \quad \text{(7 marks)}$$

Turn over for the next question

Turn over ►

- 6 (a) (i) By applying De Moivre's theorem to $(\cos \theta + i \sin \theta)^3$, show that

$$\cos 3\theta = \cos^3 \theta - 3 \cos \theta \sin^2 \theta \quad (3 \text{ marks})$$

- (ii) Find a similar expression for $\sin 3\theta$. (1 mark)

- (iii) Deduce that

$$\tan 3\theta = \frac{\tan^3 \theta - 3 \tan \theta}{3 \tan^2 \theta - 1} \quad (3 \text{ marks})$$

- (b) (i) Hence show that $\tan \frac{\pi}{12}$ is a root of the cubic equation

$$x^3 - 3x^2 - 3x + 1 = 0 \quad (3 \text{ marks})$$

- (ii) Find two other values of θ , where $0 < \theta < \pi$, for which $\tan \theta$ is a root of this cubic equation. (2 marks)

- (c) Hence show that

$$\tan \frac{\pi}{12} + \tan \frac{5\pi}{12} = 4 \quad (2 \text{ marks})$$

- 7 (a) Given that $y = \ln \tanh \frac{x}{2}$, where $x > 0$, show that

$$\frac{dy}{dx} = \operatorname{cosech} x \quad (6 \text{ marks})$$

- (b) A curve has equation $y = \ln \tanh \frac{x}{2}$, where $x > 0$. The length of the arc of the curve between the points where $x = 1$ and $x = 2$ is denoted by s .

- (i) Show that

$$s = \int_1^2 \coth x \, dx \quad (2 \text{ marks})$$

- (ii) Hence show that $s = \ln(2 \cosh 1)$. (4 marks)

END OF QUESTIONS