

Version 1.0



**General Certificate of Education (A-level)
January 2011**

Mathematics

MS/SS1B

(Specification 6360)

Statistics 1B

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
✓ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Mark Scheme – General Certificate of Education (A-level) Mathematics – Statistics 1B – January 2011

MS/SS1B

Q	Solution	Marks	Total	Comments
1(a)(i)	$r = 0.6$ to 0.98	B1		AWFW (≈ 0.8) If answers are not labelled, assume order is (a)(i) then (a)(ii)
(ii)	$r = -0.5$ to -0.02 Accept answers as ranges if and only if contained entirely within given ranges	B1	2	AWFW (≈ -0.3) Eg: (a)(i) 0.7 to $0.9 \Rightarrow$ B1 (a)(ii) -0.6 to $-0.4 \Rightarrow$ B0
(b)(i)	$r = 0.757$ $r = 0.75$ to 0.77 $r = 0.65$ to 0.85 or Attempt at $\sum x$ $\sum x^2$ $\sum y$ $\sum y^2$ and $\sum xy$ or Attempt at S_{xx} S_{yy} and S_{xy} Attempt at substitution into correct corresponding formula for r $r = 0.757$	B3 (B2) (B1) (M1) (m1) (A1)	3	AWRT (0.75708) AWFW AWFW 271.5 6142.97 1911.9 304650.01 and 43259.17 (all 5 attempted) 0.2825 36.5425 and 2.4325 (all 3 attempted) AWRT
(ii)	Strong/fairly strong/moderate positive (linear) correlation/relationship/association/link (but not 'trend') between Circumference/size and weight of (cricket) balls	Bdep1 B1	 2	Dependent on $0.65 < r < 0.85$ Or equivalent; must qualify strength and indicate positive Bdep0 for very strong/high/average/medium/some etc. Context; providing $0 < r < 1$
Total			7	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
2(a)(i)	$P(M \cap C) = \frac{175}{645} = \frac{35}{129} = 0.271$	B1	1	AWRT; accept either correct fraction
(ii)	$P(M) = \frac{519}{645} = \frac{173}{215} = 0.804 \text{ to } 0.805$	B1	1	AWFW; accept either correct fraction
(iii)	$P(LD) = \frac{63}{645} = \frac{21}{215} = 0.097 \text{ to } 0.098$	B1	1	AWFW; accept either correct fraction
(iv)	$P(L F) = \frac{94}{126} = \frac{47}{63}$ = 0.746	M1 A1	2	Accept $\frac{94}{645} \div \frac{126}{645}$ AWRT
(v)	$P(M L) = \frac{519 - 255}{645 - 349} = \frac{175 + 54 + 35}{193 + 63 + 40}$ $= \frac{264}{296} = \frac{132}{148} = \frac{66}{74} = \frac{33}{37}$ = 0.891 to 0.893	M1 M1 A1	3	Allow one arithmetic slip Allow one arithmetic slip Any of these implies M1 M1 AWFW
(b)	$P(L \cap L F) = \left(\frac{94}{126} \times \frac{93}{125} \right) \text{ or } \frac{8742}{15750}$ = 0.555	B1 B1	2	Or $\left(\frac{47}{63} \times \frac{93}{125} \right) \text{ or } \frac{4371}{7875} \text{ or } \frac{1457}{2625}$ AWRT
(c)	$P(L \cap C \cap (LD + O))$ $= \frac{349}{645} \times \frac{193}{644} \times \frac{63 + 40}{643}$ SC The three correct fractions identified but not multiplied \Rightarrow M1 M0 M0 A0 $\times 6 \text{ or } 3$ = 0.155 to 0.157 NB: 0.026 with no working \Rightarrow M1 only 0.026 \times 6 = 0.156 with no working \Rightarrow 4 marks	M1 M1 M1 A1	4	Correct numerator Correct denominator Note that a denominator of $\binom{645}{3}$ \Rightarrow M2 (second and third M1 marks) AWFW
Total			14	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	$\frac{0.98+1.00}{2}$ or $\frac{0.975+1.005}{2}$ or $0.98 + \frac{0.02}{2}$ or $0.975 + \frac{0.03}{2} = 0.99$	B1		AG (At least) one correct expression seen Ignore contradictions Accept any valid equivalent
(ii)	$\frac{0.97+0.98}{2} = 0.975$ and $\frac{1.00+1.01}{2} = 1.005$ SC In (a)(i) and (a)(ii) allow 1.0049 or 1.0049... etc	B1	2	Both CAO Can not be implied from (a)(i) Similar forms for lower boundary
(b)	Mean, $\bar{x} = 1.062$ Standard deviation, s or $\sigma = 0.043$	B1 B2	3	CAO $\sum fx = 106.2$ Ignore notation AWRT $\sum fx^2 = 112.9662$ If B0 B0, M1 can be awarded for attempt at $\frac{\sum fx}{100}$
(c)(i)	99% (0.99) $\Rightarrow z = 2.57$ to 2.58 CI for μ is $\bar{x} \pm (z \text{ or } t) \times \frac{(s \text{ or } \sigma)}{\sqrt{n}}$ Thus $1.062 \pm 2.5758 \times \frac{0.043}{\sqrt{100 \text{ or } 99}}$ Hence 1.06 ± 0.01 or (1.05, 1.07)	B1 (B1) M1 A1F A1		AWFW (2.5758) $t_{99}(0.995) = 2.626$ AWRT Used Must have \sqrt{n} with $n > 1$ F on \bar{x} , s/σ and z/t AWRT; award even if previous inaccuracies in \bar{x} , s/σ or z/t Dependent on A1F
(ii)	Volumes/ X / (parent) population may be modelled by a normal distribution / is normally distributed (Ignore contradictions)	B1	1	Or equivalent; not distribution, data, values (in table), sample, n large, nor simply 'It is stated in question'
(iii)	Sample data grouped Exact sample values unknown / mid-points used \bar{x} and s calculated from grouped data	B1	1	σ unknown s calculated from a sample \bar{x} (not μ) and s are estimates NOT data values rounded
(d)(i)	CI for μ or CI in (c)(i) > 1 LCL of CI for μ or LCL of CI in (c)(i) > 1	B1		Or equivalent; must compare CI to 1 Dependent on CI in (c)(i) > 1
(ii)	99 or 100 or all sample/ table/ data volumes/ values/ x -values/ cartons are within this range (or none/0 or 1 volumes outside)	B1	2	
	Total		13	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
4(a)	$R \sim B(15, 0.45)$			
(i)	$P(R \leq 5) = 0.26(0)$ to 0.261	B1	1	AWFW (0.2608)
(ii)	$P(R > 10) = 1 - P(R \leq 10)$ $= 1 - (0.9745 \text{ or } 0.9231)$ $= 0.025$ to 0.026	M1 A1	2	Requires '1 -' Accept 3dp rounding or truncation Can be implied by 0.025 to 0.026 but not by 0.0769 to 0.077 AWFW (0.0255)
(iii)	$P(R = 6) = 0.4522 - (a)(i)$ or $= \binom{15}{6} (0.45)^6 (0.55)^9$ $= 0.191$ to 0.192	M1 A1	2	Can be implied by a correct answer AWFW (0.1914)
(iv)	$P(5 \leq R \leq 10) = 0.9745$ or 0.9231 (p_1) Minus 0.1204 or 0.2608 (p_2) $= 0.853$ to 0.855 Or B (15, 0.45) terms stated for at least 3 values within $4 \leq R \leq 11$ gives probability $= 0.853$ to 0.855	M1 M1 A1 (M1) (A2)	3	Accept 3dp rounding or truncation $p_2 - p_1 \Rightarrow$ M0 M0 A0 $(1 - p_2) - p_1 \Rightarrow$ M0 M0 A0 $p_1 - (1 - p_2) \Rightarrow$ M1 M0 A0 only providing result > 0 Accept 3dp rounding or truncation AWFW (0.8541) Can be implied by a correct answer AWFW (0.8541)
(b)(i)	$P(S) = 0.85$ plus 1 minus (0.15×0.80) (0.15×0.20) $= 0.97$ NB: $(0.85 \times 0.20) + 0.80 \Rightarrow$ B0 B0 $(0.85 \times 0.20) + (0.85 \times 0.80)$ $+ (0.15 \times 0.80) \Rightarrow$ B0 B1	B1 B1	2	CAO; requires 'plus' or 'minus' CAO; not simply 0.12 or 0.03 AG
(ii)	$P(S \geq 48) = 0.81$ to 0.82 or 0.5553 or 0.9372 $= 0.81(0)$ to 0.811 NB: Answer = 0.4447 or 0.1892 or $0.0628 \Rightarrow$ M1 only	M2 A1	3	Accept 3dp rounding or truncation M2 for the three correctly expressed terms for B (50, 0.03) or B (50, 0.97) added AWFW (0.8108)
(iii)	$p = 1 - 0.85 = 0.15$ Mean, $\mu = 80 \times 0.15 = 12$ SC Mean = $9.6 \Rightarrow$ B1 only	B1 B1	2	CAO; may be implied by correct answer or correct expression for mean CAO
Total			15	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
5(a)	Time taken is dependent upon leaving time	B1	1	Or equivalent
(b)	b (gradient) = 1.28 (or 141/110) b (gradient) = 1.25 to 1.35 a (intercept) = 29.95 to 30 (or 659/22) a (intercept) = 29 to 31 Thus $y = 30 + 1.28x$ or Attempt at $\sum x$ $\sum x^2$ $\sum y$ and $\sum xy$ ($\sum y^2$) or Attempt at S_{xx} and S_{xy} (S_{yy}) Attempt at correct formula for b gradient b (gradient) = 1.28 (or 141/110) a (intercept) = 29.95 to 30 (or 659/22) Thus $y = 30 + 1.28x$ Accept a and b interchanged only if identified correctly by a clearly shown equation	B2 (B1) B2 (B1) B1F (M1) (m1) (A1) (A1) (B1F)	5	AWRT; (CAO or equivalent) (1.28182) AWFW Treat rounding of correct answers as ISW AWFW; (CAO or equivalent) (29.95455) AWFW F on a and b 275 9625 682 and 20575 (47494) (All four attempted) 2750 and 3525 (5210) (Both attempted) AWRT; (CAO or equivalent) AWFW; (CAO or equivalent) F on a and b If a and b are not identified anywhere in the question, then: 1.25 to 1.35 \Rightarrow B1 29 to 30 \Rightarrow B1
(c)	7.45 am $\Rightarrow x = 15$ $\Rightarrow y_{15} = 30 + 1.28 \times 15$ $= 47$ to 52 Time before 9.00 am = $9.00 - (7.45 + c's y_{15})$ $= 23$ to 28 SC Answer of 17 CAO (use of c's $y_{15} = 58$) gains 2 marks	B1 M1 A1 M1 A1	5	CAO; stated, used or implied Use of $10 < x < 20$ AWFW (49.2) May be implied AWFW (25.8) NB: An answer of 8.32 to 8.37 gains B1 M1 A1 M0 A0
(d)(i)	$y_{85} = 30 + 1.28 \times 85 = 135$ to 146	B1	1	AWFW (138.9)
(ii)	Extrapolation/ outside/ above range of x -values Implies leaves home at 8.55 so different traffic conditions	B1 B1	2	Or equivalent Or equivalent; 8.55 may be implied by 5 minutes
Total			14	

Mark Scheme – General Certificate of Education (A-level) Mathematics – Statistics 1B – January 2011

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
6(a)	Volume, $V \sim N(412, 8^2)$			
(i)	$P(V < 400) = P\left(Z < \frac{400 - 412}{8}\right)$ $= P(Z < -1.5) = 1 - P(Z < 1.5)$ $= 1 - 0.93319 = 0.066$ to 0.067	M1 M1 A1	3	Standardising 400 with 412 and 8 and/or $(412 - x)$ Area change May be implied by a correct answer or an answer < 0.5 AWFW (0.06681)
(ii)	$P(V > 420) = P(Z > 1)$ $= 1 - P(Z < 1) = 1 - 0.84134$ $= 0.158$ to 0.159	B1 B1	2	CAO but ignore inequality and sign May be implied by a correct answer AWFW (0.15866)
(iii)	$P(V = 410) = 0$ or zero or impossible	B1	1	Ignore any working B0 for 'impossible to calculate' or 'no answer'
(b)(i)	A statement/indication that (-) 1.6449 and/or 2.3263 are z-values Do not allow $\Phi(0.99) = 2.3263$, etc but allow $\Phi^{-1}(0.99) = 2.3263$ Do not award for z-value(s) simply embedded in standardisation statement(s) A clear use of $z = \frac{v - \mu}{\sigma}$ or $v = \mu + z\sigma$ with 400 and/or 420 (condone sign errors) The two given equations correctly derived	B1 M1 A1	3	Simple statement that $z = \pm 1.6449$ and/or $z = \pm 2.3263$ or sketch of normal curve with at least one z-value marked SC Immediate algebraic use of $v - \mu = z\sigma \Rightarrow$ B1 M1 A0 AG; watch for sign inconsistencies
(ii)	Thus $20 = (2.3263 + 1.6449)\sigma$ $\sigma = 5.04$ $\mu = 408$	M1 A1 A1	3	A sensible (one that would lead to values required if completed correctly) attempt at solving the two given equations by eliminating μ or σ Do NOT allow MC or MR AWRT (5.03626) AWRT (408.284)
	Total		12	
	TOTAL		75	