

OXFORD CAMBRIDGE AND RSA EXAMINATIONS**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education****MATHEMATICS****4727**

Further Pure Mathematics 3

Thursday **15 JUNE 2006** Afternoon 1 hour 30 minutes

Additional materials:
8 page answer booklet
Graph paper
List of Formulae (MF1)

TIME 1 hour 30 minutes**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

This question paper consists of 3 printed pages and 1 blank page.

2

- 1 (a) For the infinite group of non-zero complex numbers under multiplication, state the identity element and the inverse of $1 + 2i$, giving your answers in the form $a + ib$. [3]

- (b) For the group of matrices of the form $\begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}$ under matrix addition, where $a \in \mathbb{R}$, state the identity element and the inverse of $\begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix}$. [2]

- 2 (a) Given that $z_1 = 2e^{\frac{1}{6}\pi i}$ and $z_2 = 3e^{\frac{4}{3}\pi i}$, express $z_1 z_2$ and $\frac{z_1}{z_2}$ in the form $re^{i\theta}$, where $r > 0$ and $0 \leq \theta < 2\pi$. [4]

- (b) Given that $w = 2(\cos \frac{1}{8}\pi + i \sin \frac{1}{8}\pi)$, express w^{-5} in the form $r(\cos \theta + i \sin \theta)$, where $r > 0$ and $0 \leq \theta < 2\pi$. [3]

- 3 Find the perpendicular distance from the point with position vector $12\mathbf{i} + 5\mathbf{j} + 3\mathbf{k}$ to the line with equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + 5\mathbf{k} + t(8\mathbf{i} + 3\mathbf{j} - 6\mathbf{k})$. [6]

- 4 Find the solution of the differential equation

$$\frac{dy}{dx} - \frac{x^2 y}{1 + x^3} = x^2$$

for which $y = 1$ when $x = 0$, expressing your answer in the form $y = f(x)$. [8]

- 5 A line l_1 has equation $\frac{x}{2} = \frac{y+4}{3} = \frac{z+9}{5}$.

- (i) Find the cartesian equation of the plane which is parallel to l_1 and which contains the points $(2, 1, 5)$ and $(0, -1, 5)$. [5]

- (ii) Write down the position vector of a point on l_1 with parameter t . [1]

- (iii) Hence, or otherwise, find an equation of the line l_2 which intersects l_1 at right angles and which passes through the point $(-5, 3, 4)$. Give your answer in the form $\frac{x-a}{p} = \frac{y-b}{q} = \frac{z-c}{r}$. [4]

- 6 (i) Find the general solution of the differential equation

$$\frac{d^2 y}{dx^2} + 4y = \sin x. \quad [6]$$

- (ii) Find the solution of the differential equation for which $y = 0$ and $\frac{dy}{dx} = \frac{4}{3}$ when $x = 0$. [4]

3

7 The series C and S are defined for $0 < \theta < \pi$ by

$$C = 1 + \cos \theta + \cos 2\theta + \cos 3\theta + \cos 4\theta + \cos 5\theta,$$

$$S = \sin \theta + \sin 2\theta + \sin 3\theta + \sin 4\theta + \sin 5\theta.$$

(i) Show that $C + iS = \frac{e^{3i\theta} - e^{-3i\theta}}{e^{\frac{1}{2}i\theta} - e^{-\frac{1}{2}i\theta}} e^{\frac{5}{2}i\theta}$. [4]

(ii) Deduce that $C = \sin 3\theta \cos \frac{5}{2}\theta \operatorname{cosec} \frac{1}{2}\theta$ and write down the corresponding expression for S . [4]

(iii) Hence find the values of θ , in the range $0 < \theta < \pi$, for which $C = S$. [4]

8 A group D of order 10 is generated by the elements a and r , with the properties $a^2 = e$, $r^5 = e$ and $r^4a = ar$, where e is the identity. Part of the operation table is shown below.

	e	a	r	r^2	r^3	r^4	ar	ar^2	ar^3	ar^4
e	e	a	r	r^2	r^3	r^4	ar	ar^2	ar^3	ar^4
a	a	e	ar	ar^2	ar^3	ar^4				
r	r		r^2	r^3	r^4	e				
r^2	r^2		r^3	r^4	e	r				
r^3	r^3		r^4	e	r	r^2				
r^4	r^4	ar	e	r	r^2	r^3				
ar	ar		ar^2	ar^3	ar^4	a				
ar^2	ar^2		ar^3	ar^4	a	ar				
ar^3	ar^3		ar^4	a	ar	ar^2				
ar^4	ar^4		a	ar	ar^2	ar^3				

E

(i) Give a reason why D is not commutative. [1]

(ii) Write down the orders of any possible proper subgroups of D . [2]

(iii) List the elements of a proper subgroup which contains

(a) the element a , [1]

(b) the element r . [1]

(iv) Determine the order of each of the elements r^3 , ar and ar^2 . [4]

(v) Copy and complete the section of the table marked **E**, showing the products of the elements ar , ar^2 , ar^3 and ar^4 . [5]

4

BLANK PAGE