


1. (a) $(r+1)^3 - (r-1)^3 = (r^3 + 3r^2 + 3r + 1) - (r^3 - 3r^2 + 3r - 1)$
 $= \underline{6r^2 + 2}$ M1
A1 2
- (b) $\sum_{r=1}^n (6r^2 + 2) = 2^3 - 0^3$ (attempt to use an identity) M1
 $= 3^3 - 1^3$
 $4^3 - 2^3$
 \cdot
 \cdot
 \cdot
 $(n-1)^3 - (n-3)^3$
 $n^3 - (n-2)^3$
 $(n+1)^3 - (n-1)^3$ differences (must see) M1
 $= (n+1)^3 + n^3 - 1^3$ A1
- $6\sum_{r=1}^n r^2 = (n+1)^3 + n^3 - 1 - \underline{2n}$ $2n$ or equiv. B1
 $= 2n^3 + 3n^2 + n$
- $\sum_{r=1}^n r^2 = \frac{1}{6}n(2n+1)(n+1)$ (*) Sub. $\Sigma 2$ and $\div 6$ or equiv. c.s.o. M1, A1 6
[8]
2. (a) IF = $e^{\int 1 + \frac{3}{x} dx}$ M1
 $= e^{x+3\ln x}$ A1
 $= e^x e^{\ln x^3}$ must see
 $= \underline{x^3 e^x}$ A1 3
- (b) $x^3 e^x y = \int x^3 e^x \frac{1}{x^2} dx$ M1
 $= \int x e^x$
 $= x e^x - e^x + c$ \int by parts M1 A1
 $y = \frac{1}{x^2} - \frac{1}{x^3} + \frac{c}{x^3} e^{-x}$ o.e. A1 4
- (c) $I = ce^{-1} \therefore c = e^1$ M1
 $y = \frac{1}{4} - \frac{1}{8} + \frac{e \cdot e^{-2}}{8}$ M1
 $= \frac{1}{8}(1 + e^{-1})$
or = 0.171 (0.171 or better) A1 3



3. (a)

Line crosses axes
 Curve shape
 Axes contacts 6, 8, 3
 Cusps at 2 and 4

B1
 B1
 B1
 B1 4


(b) $6 - 2x = (x - 2)(x - 4)$ and $-6 + 2x = (x - 2)(x - 4)$
 $x^2 - 4x + 2 = 0$ $x^2 - 8x + 14 = 0$ either
 $x = \frac{4 \pm \sqrt{16 - 8}}{2}$ $x = \frac{8 \pm \sqrt{64 - 56}}{2}$
 $= 2 - \sqrt{2}$ $= 4 - \sqrt{2}$

M1, M1
 M1
 A1, A1 5

(c) $2 - \sqrt{2} < x < 4 - \sqrt{2}$

M1, A1 2
 [11]

4. (a) $m^2 + 4m \pm \sqrt{5-4}$ M1
 $m = \frac{-2 \pm i}{2}$
- $= \frac{-2 \pm i}{2}$ A1
 $y = e^{-2x}(A \cos x \pm B \sin x)$ M1
- PI = $\lambda \sin 2x + \mu \cos 2x$ PI & attempt diff. M1
 $y' = 2\lambda \cos 2x - 2\mu \sin 2x$
 $y'' = -4\lambda \sin 2x - 4\mu \cos 2x$ A1
 $\therefore -4\lambda - 8\mu + 5\lambda = 65$
 $-4\mu + 8\lambda + 5\mu = 0$ subst. in eqn. & equate M1
 $\lambda - 8\mu = 65$
 $8\lambda + \mu = 0$ solving sim. eqn. M1
 $64\lambda + 8\mu = 0$
 $65\lambda = 65$
 $\lambda = 1, \mu = -8$ A1
- $\therefore y = e^{-2x}(A \cos x + B \sin x) + \sin 2x - 8 \cos 2x$ on their λ and μ A1ft 9
- (b) As $x \rightarrow \infty, e^{-2x} \rightarrow 0 \therefore y \rightarrow \sin 2x - 8 \cos 2x$ B1ft
 $y \rightarrow R \sin(2x + \alpha)$ M1
 $R = \sqrt{65}$
 $\alpha = \tan^{-1} -8 = -1.446$ or -82.9° A1 3
[12]



axis

Shape + horiz.

B1

3

B1

2

(b) Area = $\frac{1}{2} \int r^2 d\theta$
= $\frac{1}{2} \int \frac{9 \cos^2 2\theta + 1}{2} d\theta$ use of $\frac{1}{2} \int r^2$ M1
= $\frac{9}{2} \int \frac{\cos 4\theta + 1}{2} d\theta$ use of $\cos 4\theta = 2\cos^2 2\theta - 1$ M1
= $\frac{9}{2} \left[\frac{\sin 4\theta}{8} + \frac{\theta}{2} \right]_{\frac{\pi}{6}}^{\frac{\pi}{4}}$
 \int
= $\frac{9}{2} \left[\frac{\pi}{8} - \frac{\sqrt{3}}{16} - \frac{\pi}{12} \right]$ M1, A1
= $\frac{9}{2} \left[\frac{\pi}{24} - \frac{\sqrt{3}}{16} \right]$ subst. $\frac{\pi}{4}$ and $\frac{\pi}{6}$ M1
= $2 \left[\frac{\pi}{24} - \frac{\sqrt{3}}{16} \right]$ or 0.103 A1 6

(c) $r \sin \theta = 3 \sin \theta \cos 2\theta$
 $\frac{d'y'}{d\theta} = 3 \cos \theta \cos 2\theta - 6 \sin \theta \sin 2\theta$ (diff. $r \sin \theta$) M1, A1
 $\frac{dy}{d\theta} = 0 \Rightarrow 6 \cos^2 \theta - 3 \cos \theta - 12 \sin^2 \theta \cos \theta = 0$ use of $\frac{dy}{d\theta} = 0$ M1
 $6 \cos^2 \theta - 3 \cos \theta - 12(1 - \cos^2 \theta) \cos \theta = 0$ use double angle formula M1
 $18 \cos^3 \theta - 15 \cos \theta = 0$ solving M1
 $\cos \theta = 0$ or $\cos^2 \theta = \frac{5}{6}$ or $\tan^2 \theta = \frac{1}{5}$ or $\sin^2 \theta = \frac{1}{6}$ A1
 $\therefore r = 3(2 \times \frac{5}{6}) - 1$
= 2
 $\therefore r \sin \theta = 2 \sqrt{\frac{1}{6}}$ use of $d = 2r \sin \theta$ M1
 $\Rightarrow d = \frac{2\sqrt{6}}{3}$ A1 8
[16]

6. Solves $x^2 - 2 = 2x$ by valid method M1
Obtains $x = 1 \pm \sqrt{3}$ or equivalent A1
(may only obtain relevant root if graph is used)
Solves $2 - x^2 = 2x$ M1
Obtains $x = -1 \pm \sqrt{3}$ A1
Rejects two of these roots and obtains (or uses graph and obtains) dM1
 $x > 1 + \sqrt{3}, x < -1 + \sqrt{3}$ A1, A1 7

Special case:

Squares both sides to obtain quadratic in x^2 and solve to obtain $x^2 = 4 \pm 2\sqrt{3}$

Obtains $x = 1 \pm \sqrt{3}$ or $x = -1 \pm \sqrt{3}$

Last three marks as before.

M1A1

M1A1

dM1A1A1

[7]

7. (a) Integrating Factor = e^{2x}

B1

$$\frac{d}{dx}(ye^{2x}) = xe^{2x}$$

M1

$$ye^{2x} = \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x} dx$$

M1

$$= \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + c$$

A1

Min point and passing through (0, 1)

$$\therefore y = \frac{1}{2}x - \frac{1}{4} + ce^{-2x}$$

A1

5

shape

(b) $1 = c - \frac{1}{4} \rightarrow c = \frac{5}{4}$

M1

$$\therefore y = \frac{1}{2}x - \frac{1}{4} + \frac{5}{4}e^{-2x} \text{ and } \frac{d}{dx} = \frac{1}{2} - \frac{5}{2}e^{-2x}$$

M1

When $y' = 0$, $e^{-2x} = \frac{1}{5} \therefore 2x = \ln 5$


M1

$x = \frac{1}{2} \ln 5$, $y = \frac{1}{4} \ln 5$ at minimum point.

A1

4

(c)



B1B1 2
[11]

8. (a) Auxiliary equation: $m^2 + 2m + 2 = 0 \rightarrow m = -1 \pm i$
Complementary Function is $y = e^{-1} (A \cos t + B \sin t)$
Particular Integral is $y = \lambda e^{-1}$, with $y' = -\lambda e^{-1}$, and $y'' = \lambda e^{-1}$
 $\therefore (\lambda - 2\lambda + 2\lambda)e^{-1} = 2e^{-1} \rightarrow \lambda = 2$
 $\therefore y = e^{-1}(A \cos t + B \sin t + 2)$

M1
M1A1
M1
A1
B1 6

- (b) Puts $1 = A + 2$ and solves to obtain $A = -1$
 $y' = e^{-1}(-A \sin t + B \cos t) - e^{-1}(A \cos t + B \sin t + 2)$
Puts $1 = B - A - 2$ and uses value for A to obtain B
 $B = 2$
 $\therefore y = e^{-1}(2 \sin t - \cos t + 2)$

M1 A1ft
M1
A1cso
A1cso 6
[12]

9. (a) $3a(1 - \cos \theta) = a(1 + \cos \theta)$
 $2a = 4a \cos \theta \rightarrow \cos \theta = \frac{1}{2} \therefore \theta = \frac{\pi}{3} \text{ or } -\frac{\pi}{3}$
 $r = \frac{3a}{2}$

M1
M1
A1A1 4

[Co-ordinates of points are $(\frac{3a}{2}, \frac{\pi}{3})$ and $(\frac{3a}{2}, -\frac{\pi}{3})$]

(b) $AB = 2r\sin\theta = \frac{3a\sqrt{3}}{2}$ M1A1 2

$$\text{Area} = \int_{-\frac{\pi}{3}}^{\frac{\pi}{3}} \frac{1}{2} r^2 d\theta$$

$$= \frac{1}{2} \int [a^2(1 + \cos\theta)^2 - 9a^2(1 - \cos\theta)^2] d\theta$$

M1 M1

$$= \frac{a^2}{2} \int [1 + 2\cos\theta + \cos^2\theta - 9(1 - 2\cos\theta + \cos^2\theta)] d\theta$$

A1

$$= \frac{a^2}{2} \int [-8 + 20\cos\theta - 8\cos^2\theta] d\theta$$

$$= k[-8\theta + 20\sin\theta \dots$$

B1

$$\dots - 2\sin 2\theta - 4\theta]$$

B1

Uses limits $\frac{\pi}{3}$ and $-\frac{\pi}{3}$ correctly or uses twice smaller area

and uses limits $\frac{\pi}{3}$ and 0 correctly. (Need not see 0 substituted)

$$= a^2[-4\pi + 10\sqrt{3} - \sqrt{3}] \text{ or } = a^2[-4\pi + 9\sqrt{3}] \text{ or } 3.022a^2$$

A1 7

(d) $3a\frac{\sqrt{3}}{2} = 4.5 \rightarrow a = \sqrt{3}$ B1

$$\therefore \text{Area} = 3[9\sqrt{3} - 4\pi], = 9.07 \text{ cm}^2$$

M1, A1 3
[16]

10. (a) $f'(x) = \sec^2 x$ $f''(x) = 2\sec x(\sec x \tan x)$ (or equiv.) M1 A1

$$f''(x) = 2\sec^2 x(\sec^2 x) + 2 \tan x(2\sec^2 x \tan x)$$
 (or equiv.) A1 3

$$(2\sec^2 x + 6\sec^2 x \tan^2 x)$$

$$(2\sec^4 x + 4\sec^2 x \tan^2 x), (6\sec^4 x - 4\sec^2 x), (2 + 8\tan^2 x + 6\tan^4 x)$$

(b) $\tan\frac{\pi}{4} = 1$ or $\sec\frac{\pi}{4} = \sqrt{2}$ (1, 2, 4, 16) B1

$$\tan x = f\left(\frac{\pi}{4}\right) + \left(x - \frac{\pi}{4}\right)f'\left(\frac{\pi}{4}\right) + \frac{1}{2}\left(x - \frac{\pi}{4}\right)^2 f''\left(\frac{\pi}{4}\right) + \frac{1}{6}\left(x - \frac{\pi}{4}\right)^3 f'''\left(\frac{\pi}{4}\right)$$

M1

$$= 1 + 2\left(x - \frac{\pi}{4}\right) + 2\left(x - \frac{\pi}{4}\right)^2 + \frac{8}{3}\left(x - \frac{\pi}{4}\right)^3$$

A1(cso)3

(Allow equiv. fractions)

$$(c) \quad x = \frac{3\pi}{10}, \text{ so use } \left(\frac{3\pi}{10} - \frac{\pi}{4} \right) \left(= \frac{\pi}{20} \right)$$

$$\tan \frac{3\pi}{10} \approx 1 + \frac{\pi}{10} + \left(2 \times \frac{\pi^2}{400} \right) + \left(\frac{8}{3} \times \frac{\pi^3}{8000} \right) = 1 + \frac{\pi}{10} + \frac{\pi^2}{200} + \frac{\pi^3}{3000} \quad (*)$$

M1

A1(cso)2

[8]

11. (a) $n = 1: \frac{d}{dx} (e^x \cos x) = e^x \cos x - e^x \sin x$ M1

(Use of product rule)

$$\cos \left(x + \frac{\pi}{4} \right) = \cos x \cos \frac{\pi}{4} - \sin x \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}} (\cos x - \sin x)$$
 M1

$$\frac{d}{dx} (e^x \cos x) = 2^{1/2} e^x \cos \left(x + \frac{\pi}{4} \right) \quad \text{True for } n = 1 \text{ (c.s.o. + comment)}$$
 A1

Suppose true for $n = k$.

$$\left[\frac{d^{k+1}}{dx^{k+1}} (e^x \cos x) \right] = \frac{d}{dx} \left(2^{1/2} e^x \cos \left(x + \frac{k\pi}{4} \right) \right)$$
 M1

$$= 2^{1/2} \left[e^x \cos \left(x + \frac{k\pi}{4} \right) - e^x \sin \left(x + \frac{k\pi}{4} \right) \right]$$
 A1

$$= 2^{1/2} e^x \sqrt{2} \cos \left(x + \frac{k\pi}{4} + \frac{\pi}{4} \right) = 2^{1/2(k+1)} e^x \cos \left(x + (k+1) \frac{\pi}{4} \right)$$
 M1 A1

\therefore True for $n = k + 1$, so true (by induction) for all n . (≥ 1) A1(cso)8

(b) $1 + \left(\sqrt{2} \cos \frac{\pi}{4} \right) x + \frac{1}{2} \left(2 \cos \frac{\pi}{2} \right) x^2 + \frac{1}{6} \left(2\sqrt{2} \cos \frac{3\pi}{4} \right) x^3 + \frac{1}{24} (4 \cos \pi) x^4$ M1

(1) (0) (-2) (-4)

$$e^x \cos x = 1 + x - \frac{1}{3} x^3 - \frac{1}{6} x^4 \quad \text{(or equiv. fractions)}$$
 A2(1,0)3

[11]


12. (a) $\arg z = \frac{\pi}{4} \Rightarrow z = \lambda + \lambda i$ (or putting x and y equal at some stage) B1

$$w = \frac{(\lambda + 1) + \lambda i}{\lambda + (\lambda + 1)i}, \text{ and attempt modulus of numerator or denominator.}$$
 M1

(Could still be in terms of x and y)

$$|(\lambda + 1) + \lambda i| = |\lambda + (\lambda + 1)i| = \sqrt{(\lambda + 1)^2 + \lambda^2}, \therefore |w| = 1 \quad (*)$$
 A1, A1cso

- (b) $w = \frac{z+1}{z+i} \Rightarrow zw + wi = z + 1 \Rightarrow z = \frac{1-wi}{w-1}$ M1
 $|z| = 1 \Rightarrow |1-wi| = |w-1|$ M1 A1
 For $w = \frac{a+ib}{1+b}$, $|(1+ib)\frac{a+ib}{1+b}| = |(a-1)+ib|$ M1
 $\sqrt{(1+b)^2 + a^2} = \sqrt{(a-1)^2 + b^2}$ M1
 $b = -a$ A1 Image is (line) $y = -x$ 6

- (c)  B1 B1 2

- (d) $z = i$ marked (P) on z -plane sketch. B1
 $z = i \Rightarrow \frac{1+i}{2i} = \frac{i-1}{-2} = \frac{1}{2} - \frac{1}{2}i$ marked (Q) on w -plane sketch. B1 2

[14]