



GCE

Mathematics

Advanced GCE

Unit 4724: Core Mathematics 4

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

- 1 Attempt to factorise **both** numerator & denominator M1 completely or partially
 Num = e.g. $(x^2 - 1)(x^2 - 9)$ or $(x^2 - 2x - 3)(x^2 + 2x - 3)$ B1 or $(x - 3)(x + 3)(x - 1)(x + 1)$
 Denominator = e.g. $(x^2 - 2x - 3)(x + 5)(x + 3)$ B1 or $(x - 3)(x + 1)(x + 5)(x + 3)$
 $\frac{x-1}{x+5}$ or $1 - \frac{6}{x+5}$ WWW A1 4 ISW but not if any further 'cancellation'

Alternative start, attempting long division

Expand denom as quartic & attempt to divide $\frac{\text{numerator}}{\text{denominator}}$ M1 but not divide $\frac{\text{denominator}}{\text{numerator}}$

Obtain quotient = 1 & remainder = $-6x^3 - 6x^2 + 54x + 54$ B1

Final B1 A1 available as before

4

- 2 $2^2 + (-3)^2 + (\sqrt{12})^2$ soi e.g. 25 or 5 M1 Allow $2^2 - 3^2 + \sqrt{12}^2$
 5 A1 May be implied by 5 or 1/5 in final answer

$\frac{1}{5} \begin{pmatrix} 2 \\ -3 \\ \sqrt{12} \end{pmatrix}$ or $\begin{pmatrix} \frac{2}{5} \\ -\frac{3}{5} \\ \frac{\sqrt{12}}{5} \end{pmatrix}$ AEF $\sqrt{A1}$ 3 FT their '5'. Accept $-\frac{1}{5} \begin{pmatrix} \\ \\ \phantom{\sqrt{12}} \end{pmatrix}$ or $\frac{1}{\pm 5} \begin{pmatrix} \\ \\ \phantom{\sqrt{12}} \end{pmatrix}$

3

- 3 (i) The words quotient and remainder need not be explicit

Long division For leading term $3x$ in quotient B1

Suff evidence of div process ($3x$, mult back, attempt sub) M1

(Quotient) = $3x - 1$ A1

(Remainder) = x **AG** A1 4 No wrong working, partic on penult line

Identity $3x^3 - x^2 + 10x - 3 = Q(x^2 + 3) + R$ *M1

$Q = ax + b, R = cx + d$ & attempt at least 2 operations dep*M1 If $a = 3$, this \Rightarrow 1 operation

$a = 3, b = -1$ A1

$c = 1, d = 0$ A1 No wrong working anywhere

Inspection $3x^3 - x^2 + 10x - 3 = (x^2 + 3)(3x - 1) + x$ B2 or state quotient = $3x - 1$

Clear demonstration of LHS = RHS B2

- (ii) Change integrand to 'their (i) quotient' + $\frac{x}{x^2 + 3}$ M1

Correct FT integration of 'their (i) quotient' $\sqrt{A1}$

$\int \frac{x}{x^2 + 3} dx = \frac{1}{2} \ln(x^2 + 3)$ A1

Exact value of integral = $\frac{1}{2} + \frac{1}{2} \ln 4 - \frac{1}{2} \ln 3$ AEF ISW A1 4 Answer as decimal value (only) \rightarrow A0

8

- 4 Indefinite integral Attempt to connect dx and $d\theta$ M1 Incl $\frac{dx}{d\theta} = \frac{d\theta}{dx} = \dots$, $dx = \dots d\theta$; not $dx = d\theta$
- Denominator $(1-9x^2)^{\frac{3}{2}}$ becomes $\cos^3\theta$ B1
- Reduce original integral to $\frac{1}{3} \int \frac{1}{\cos^2\theta} d\theta$ A1 May be implied, seen only as $\frac{1}{3} \int \sec^2\theta d\theta$
- Change $\int \frac{1}{\cos^2\theta} d\theta$ to $\tan\theta$ B1 Ignore $\frac{1}{3}$ at this stage
- Use appropriate limits for θ (allow degrees) or x M1 Integration need not be accurate
- $\frac{\sqrt{3}}{9}$ AEF, exact answer required, ISW A1 **6**

6

- 5 (i) Attempt to set up 3 equations M1 of type $4 + 3s = 1, 6 + 2s = t, 4 + s = -t$
- $(s, t) = (-1, 4)$ or $(-1, -3)$ or $(-\frac{10}{3}, -\frac{2}{3})$ *A1 or $s = -1$ & $-\frac{10}{3}$ or $t =$ two of $(4, -3, -\frac{2}{3})$
- Show clear contradiction e.g. $3 \neq -4, 4 \neq -3, -6 \neq 1$ dep*A1 **3** Allow \checkmark unsimpl contradictions. No ISW.
- SC If $s = -\frac{10}{3}$ found from 2nd & 3rd eqns and contradiction shown in 1st eqn, all 3 marks may be awarded.

- (ii) Work with $\begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$ M1

Clear method for scalar product of any 2 vectors M1

Clear method for modulus of any vector M1

79.1^(e) or better (79.1066..) 1.38 (rad) (1.38067..) ISW A1 **4** (From $\frac{1}{\sqrt{14} \cdot \sqrt{2}}$)

- (iii) Use $\begin{pmatrix} 4+3s \\ 6+2s \\ 4+s \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} = 0$ M1

Obtain $s = -2$ A1 from $12 + 9s + 12 + 4s + 4 + s = 0$

A is $\begin{pmatrix} -2 \\ 2 \\ 2 \end{pmatrix}$ or $-2\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ final answer B1 **3** Accept $(-2, 2, 2)$

10

6 $(1+ax)^{1/2} = 1 + \frac{1}{2}ax + \dots + \frac{\frac{1}{2} \cdot \frac{-1}{2}}{2} (ax)^2$ B1, B1 N.B. third term = $-\frac{1}{8}a^2x^2$

Change $(4-x)^{-1/2}$ into $k(1-\frac{x}{4})^{-1/2}$, where k is likely to be $\frac{1}{2}/2/4/-2$, & work out expansion of $(1-\frac{x}{4})^{-1/2}$

$(1-\frac{x}{4})^{-1/2} = 1 + \frac{1}{8}x + \dots + \frac{\frac{-1}{2} \cdot \frac{-3}{2}}{2} (\frac{-x}{4})^2$ B1, B1 N.B. third term = $\frac{3}{128}x^2$

OR Change $\{4-x\}^{1/2}$ into $l(1-\frac{x}{4})^{1/2}$, where l is likely to be $\frac{1}{2}/2/4/-2$, & work out expansion of $(1-\frac{x}{4})^{1/2}$

$(1-\frac{x}{4})^{1/2} = 1 - \frac{1}{8}x - \frac{1}{128}x^2$ B1 (for all 3 terms simplified)

$k = \frac{1}{2}$ (with possibility of M1 + A1 + A1 to follow) B1 $l = 2$ (with no further marks available)

Multiply $(1+ax)^{1/2}$ by $(4-x)^{-1/2}$ or $(1-\frac{x}{4})^{-1/2}$ M1 Ignore irrelevant products

The required three terms (with/without x^2) identified as

$-\frac{1}{16}a^2 + \frac{1}{32}a + \frac{3}{256}$ or $\frac{-16a^2+8a+3}{256}$ AEF ISW A1+A1 8 A1 for one correct term + A1 for other two

SC B1 for $\frac{1}{4}(1-\frac{x}{4})^{-1}$; B1 for $(1-\frac{x}{4})^{-1} = 1 + \frac{x}{4} + \frac{x^2}{16}$; M1 for multiplying $(1+ax)$ by their $(4-x)^{-1}$.

If result is $p+qx+rx^2$, then to find $(p+qx+rx^2)^{1/2}$ award B1 for $p^{1/2}(\dots)$,

B1 correct 1st & 2nd terms of expansion, B1 correct 3rd term; A1, A1 as before, for correct answers.

8

7 Attempt to sep variables in format $\int py^2 (dy) = \int \frac{q}{x+2} (dx)$ M1 where constants p and/or q may be wrong

Either y^3 & $\ln(x+2)$ or $\frac{1}{3}y^3$ & $\frac{1}{3}\ln(x+2)$ A1+A1 Accept $\frac{1}{3}\ln(3x+6)$ for $\frac{1}{3}\ln(x+2)$ & $|$ for ()

If indefinite integrals are being used (most likely scenario)

Substitute $x=1, y=2$ into an eqn containing '+const' M1

Sub $y=1.5$ and their value of 'const' & solve for x or q M1

x or $q = -1.97$ only A2

[SC x or $q = -1.970$ or -1.971 or -1.9705 or -1.9706 A1] 7

If definite integrals are used (less likely scenario)

Use $\int_{1.5}^2 \dots dy = \int_q^1 \dots dx$ where 2 corresponds with 1..... M2 & 1.5 corresp with q (at top/bottom or v.v.)

Then A2 or SC A1 as above

Use $\int_{1.5}^2 \dots dy = \int_1^q \dots dx$ where 2 corresponds with q M1 & 1.5 corresp with 1 (at top/bottom or v.v.)

Then A1 for 1.97 only

7

8 Cartesian equation may be used in parts (i) - (iii) and corresponding marks awarded

(i) Sub parametric eqns into $y = 3x$ & produce $t = -2$

OR sub $t = -2$ into para eqs, obtain $(-1, -3)$ & state $y = 3x$

OR other similar methods producing (or verifying) $t = -2$ B1

Value of t at other point is 2 B1 2 $t = \pm 2$ is sufficient for B1+B1

(ii) Use (not just quote) $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$ M1

$= -(t+1)^2$ A1 or $\frac{-1}{x^2}$ or $\frac{-(2+y)}{x}$

Attempt to use $-\frac{1}{\frac{dy}{dx}}$ for gradient of normal M1

Gradient normal = 1 cao A1

Subst $t = -2$ into the parametric eqns. M1 to find pt at which normal is drawn

Produce $y = x - 2$ as equation of the normal WWW A1 6 'A' marks in (ii) are dep on prev 'A'

(iii) Substitute the parametric values into their eqn of normal M1

Produce $t = 0$ as final answer cao A1 2 This is dep on final A1 in (ii)

N.B. If $y = x - 2$ is found fortuitously in (ii) (& \therefore given A0 in (ii)), you must award A0 here in (iii).

(iv) Attempt to eliminate t from the parametric equations M1

Produce any correct equation A1 e.g. $x = \frac{1}{y+2}$

Produce $y = \frac{1}{x} - 2$ or $y = \frac{1-2x}{x}$ ISW A1 3 Must be seen in (iv)

{N.B. Candidate producing only $y = \frac{1}{x} - 2$ is awarded both A1 marks.}

4724

Mark Scheme

June 2011

- 9 (i) Treat $x \ln x$ as a product M1 If $\int \ln x$, use parts $u = \ln x, dv = 1$
- Obtain $x \cdot \frac{1}{x} + \ln x$ A1 $x \ln x - \int 1 dx = x \ln x - x$
- Show $x \cdot \frac{1}{x} + \ln x - 1 = \ln x$ WWW AG A1 3 And state given result

(ii)(a) Part (a) is mainly based on the indef integral $\int (\ln x)^2 dx$

[A candidate stating e.g. $\int (\ln x)^2 dx = \int 2 \ln x dx$ or $= \int (\ln x - x)^2 dx$ is awarded 0 for (ii)(a)]

Correct use of $\int \ln x dx = x \ln x - x$ anywhere in this part B1 Quoted from (i) or derived

Use integ by parts on $\int (\ln x)^2 dx$ with $u = \ln x, dv = \ln x$ M1 or $u = (\ln x)^2, dv = 1$

[For 'integration by parts, candidates must get to a 1st stage with format $f(x) + /- \int g(x) dx$]

1st stage = $\ln x(x \ln x - x) - \int \frac{1}{x}(x \ln x - x) dx$ soi A1 $x(\ln x)^2 - \int x \cdot \frac{2}{x} \ln x dx$

2nd stage = $x(\ln x)^2 - 2x \ln x + 2x$ AEF (unsimplified) A1

\therefore Value of definite integral between 1 & e = e - 2 cao A1 Use limits on 2nd stage & produce cao

Volume = $\pi(e - 2)$ ISW A1 6 Answer as decimal value (only) \rightarrow A0

Alternative method when subst. $u = \ln x$ used

Attempt to connect dx and du M1

Becomes $\int u^2 e^u du$ A1

First stage $u^2 e^u - \int 2u e^u du$ A1

Third stage $(u^2 - 2u + 2)e^u$ A1

Final A1 A1 available as before

(b) Indication that reqd vol = vol cylinder - vol inner solid M1

Clear demonstration of either vol of cylinder being πe^2
(including reason for height = $\ln e$) or rotation of $x = e$

about the y-axis (including upper limit of $y = \ln e$) A1 Could appear as $\pi \int_0^1 e^2 dy$

$(\pi) \int x^2 dy = (\pi) \int e^{2y} dy$ B1

$\frac{\pi(e^2 + 1)}{2}$ or 13.2 or 13.18 or better B1 4 May be from graphical calculator

13

Possible helpful points

1. M is Method; does the candidate know what he/she should be doing? It does not ask how accurate it is.. e.g. in Qu.4, a candidate saying $\frac{dx}{d\theta} = -\frac{1}{3} \cos \theta$ is awarded M1.
2. When checking if decimal places are acceptable, accept both rounding & truncation.
3. In general we ISW unless otherwise stated.
4. The symbol \surd is sometimes used to indicate 'follow-through' in this scheme.

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553