





PH2

Question		Marking details	Marks Available		
1	(a)	(i)		2	
		(ii)	Either $\lambda = 1.16 \text{ [m]} (1)$ $f = 50 \text{ [Hz]} (1)$ $v = 58 \text{ [m s}^{-1}\text{]}(1)$		Or $\lambda = 1.16 \text{ [m]} (1)$ $v = \frac{\lambda}{T} \text{ or } v = \frac{1.16}{0.02} (1)$ $v = 58 \text{ [m s}^{-1}\text{]}(1)$
		(iii)	[1 mark only if either 1.2m used or 1.74/0.03 used]		3
		(iv)	All 4 nodes labelled		1
	(b)	(i)		1	
		(ii)	Either line drawn ✓ $f = 17 \text{ Hz} (1)$ UNIT mark New wavelength = 3.48 m or 3 × previous λ or appeal to $f = [n] \frac{v}{2L} (1)$ (Allow 1 mark only if $f = 34 \text{ Hz}$) Allow e.c.f. from (b)(i)	2	
	(c)	(i)	The displacement at any point is the [vector] sum of the displacements of the individual waves.	1	
		(ii)	$t = 1.0 \text{ s}$: horizontal line shown (1) $t = 2.0 \text{ s}$: inversion of $t = 0$ shown (1)	2	
	Question 1 total			[13]	

Question		Marking details	Marks Available	
2	(a)	(i)	I. Vibrations / oscillations / displacements [accept particle displacements] are perpendicular / at right angles / 90° to the propagation directions [or equiv.]	1
			II. Vibrations / oscillations / displacements [accept particle displacements] are in one direction [accept in one plane]	1
		(ii)	Alternates [gradually] between light and dark (1)	
			2 extinctions / dark places in 360°/ or equivalent (1) [Accept an answer which assumes initially bright or initially dark]	2
	(b)	(i)	I. Light spreads out [round edge of each slit] [or equiv.]	1
			II. So light from the two slits overlaps [or equiv.]	1
		(ii)	I. $\lambda = \frac{2.0 \text{ mm} \times 0.50 \text{ mm}}{1.5 \text{ m}}$ (1) = 670 n[m] (1) [667 nm, accept 700 nm]	2
			II. Fringe separation increased (1); [bright] fringes dimmer (1)	2
	(c)	$3\lambda = d \sin 77^\circ$ [or by impl.] (1) $d = \frac{1}{5.00 \times 10^5} \text{ m} [= 2.00 \times 10^{-6} \text{ m}]$ [or by impl.] (1) $\lambda = 650 \text{ n[m]}$ (1)	3	
	Question 2 total			[13]

Question		Marking details	Marks Available										
3	(a)	(i) Smooth curve drawn through all the points (ii) 46° [or as appropriate from drawn line] (iii) Reflected ray drawn with angle of reflection equal to θ_p by eye . (iv) Any of: <table style="display: inline-table; border-collapse: collapse; vertical-align: middle;"> <tr> <td style="border-right: 1px solid black; padding: 5px;">$\frac{\sin 14^\circ}{\sin 10^\circ} \checkmark$</td> <td style="border-right: 1px solid black; padding: 5px;">$\frac{\sin 28.5^\circ}{\sin 20^\circ}$</td> <td style="border-right: 1px solid black; padding: 5px;">$\frac{\sin 44^\circ}{\sin 30^\circ}$</td> <td style="border-right: 1px solid black; padding: 5px;">$\frac{\sin 64^\circ}{\sin 40^\circ}$</td> <td style="padding: 5px;">$\frac{\sin 82^\circ}{\sin 45^\circ}$</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 5px;">$1.39 [\pm 0.05] \checkmark$</td> <td style="border-right: 1px solid black; padding: 5px;">$1.40 [\pm 0.05]$</td> <td style="border-right: 1px solid black; padding: 5px;">$1.40 [\pm 0.05]$</td> <td style="border-right: 1px solid black; padding: 5px;">$1.40 [\pm 0.05]$</td> <td style="padding: 5px;">$1.40 [\pm 0.05]$</td> </tr> </table> or by implication (v) I. Any $2 \times (1)$ from <ul style="list-style-type: none"> • Straight \checkmark • Through the origin \checkmark • Gradient $> 1 \checkmark$ II. [n is the] gradient	$\frac{\sin 14^\circ}{\sin 10^\circ} \checkmark$	$\frac{\sin 28.5^\circ}{\sin 20^\circ}$	$\frac{\sin 44^\circ}{\sin 30^\circ}$	$\frac{\sin 64^\circ}{\sin 40^\circ}$	$\frac{\sin 82^\circ}{\sin 45^\circ}$	$1.39 [\pm 0.05] \checkmark$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$	1 1 1 2 2 1
	$\frac{\sin 14^\circ}{\sin 10^\circ} \checkmark$	$\frac{\sin 28.5^\circ}{\sin 20^\circ}$	$\frac{\sin 44^\circ}{\sin 30^\circ}$	$\frac{\sin 64^\circ}{\sin 40^\circ}$	$\frac{\sin 82^\circ}{\sin 45^\circ}$								
$1.39 [\pm 0.05] \checkmark$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$	$1.40 [\pm 0.05]$									
(b)	(i) $1.530 \sin c = 1.520 [\sin 90^\circ] (1)$ [or by impl.] $c = 83^\circ (1)$ (ii) $\theta = 7^\circ$ [accept 6.5°] e.c.f. from (b)(i) (iii) Smaller <u>differences</u> in distances travelled or times taken [by light travelling different paths] (1), so less blurring / smearing / overlap of data / pulses (1) [or data can be transmitted at a greater rate] Less multimode dispersion only award 2 nd mark Question 3 Total	2 1 2 [13]											
4	(a)	$f_{\text{Thresh}} = \frac{\phi}{h} (1)$ [or by impl.] = $5.1[3] \times 10^{14}$ [Hz] (1)	2										
	(b)	(i) Photon $E = 6.63 \times 10^{-34} \times 7.4 \times 10^{14} [= 4.91 \times 10^{-19} \text{J}]$ [or by impl.] (1) $E_{k \text{ max}} [= 4.91 \times 10^{-19} - 3.4 \times 10^{-19}] = 1.5 \times 10^{-19}$ [J] (1) (ii) [A single] photon gives its energy to an electron (1) Some of the energy used to escape from the metal (1).	2 2										
(c)	(i) Points plotted at $(5.1 \times 10^{14} \text{Hz}, 0)$ and $(7.4 \times 10^{14} \text{Hz}, 1.5 \times 10^{-19} \text{J})$ (1) Allow e.c.f. from (a) and (b)(i) Straight line drawn through points (1) (One correct point only and a positive slope line = 1 mark) (ii) h / the Planck constant (iii) Straight line drawn with same gradient as (i) and to the right	2 1 1											
		Question 4 Total	[10]										

Question		Marking details	Marks Available
5	(a)	$E = \frac{hc}{\lambda}$ [or equiv. eg. $E = hf$ and $\lambda = \frac{c}{f}$ or by impl] (1) $\lambda_{UG} = 6.95 \times 10^{-7}$ [m] (1)	2
	(b)	(i) More electrons in level U than in level G	1
		(ii) They / the photons would be absorbed [accept 'disappear'] (1). The energy would be used to excite ions [accept atoms] / raise electrons from G to U [or equiv.] (1)	2
		(iii)  <p style="text-align: center;">Both transitions shown</p>	1
(c)	(iv) Any 2 x (1) from <ul style="list-style-type: none"> • Passing / incident photon ✓ • Excited ion ✓ • Electron drops to lower level ✓ • The incident photon must have wavelength = λ_{UG} [or 695 nm] or must have energy 2.86×10^{-19} J ✓ 3 rd mark <ul style="list-style-type: none"> • 2 photons where there was one previously. Accept by implication e.g. in phase with the incident photon. 	3	
		Any 2 → (1); any third →(2) from <ul style="list-style-type: none"> • [plane] polarised ✓ • Coherent ✓ • Monochromatic ✓ • Parallel beam ✓ 	2
Question 5 Total			[11]

Question		Marking details	Marks Available
6	(a)	(i) $\lambda_{\text{peak}} = \frac{2.90 \times 10^{-3} \text{ K m}}{2.5 \times 10^7 \text{ K}} (1) = 1.16 \times 10^{-10} \text{ [m]} (1)$	2
		(ii) X-ray / γ -ray	1
		(iii) 	1
		(iv) Spectral intensity low <u>in high λ 'tail'</u> but not zero.	1
	(b)	$P = \sigma A \times (2.5 \times 10^7 \text{ K})^4$ [or by impl.] (1) $A = 4\pi \times 11000^2$ [or by impl.] (1) [= $1.52 \times 10^9 \text{ m}^2$] $P = 3.4 \times 10^{31} \text{ W}$ (1) UNIT mark	3
	(c)	$A_2 T_2^4 = A_1 T_1^4$ (1) or $T_2^4 = \frac{3.4 \times 10^{31}}{5.67 \times 10^{-8} \times 3.04 \times 10^9} \text{ K}^4$ e.c.f from (b) $T_2 = 2.1 \times 10^7 \text{ K}$ (1)	2
Question 6 Total			[10]
7	(a)	(i) Any 3 \times (1) from <ul style="list-style-type: none"> • d have $\frac{1}{3}$ electronic charge / $-\frac{1}{3}e$ charge ✓ • ds have greater mass than e ✓ • ds feel strong force [or interact with gluons]; e don't ✓ • ds cannot be isolated; e can [or d can only be found in specific groupings; e can be by itself] ✓ • ds have lepton number 0, es have lepton number 1 ✓ 	3
		(ii) $[3 \times (-\frac{1}{3}e)] = -e$ [accept e or -1 or $1.6 \times 10^{-19} \text{ C}$ with some justification]	1
		(b) Any 2 \times (1) from <ul style="list-style-type: none"> • Very short decay time ✓ • Individual quark flavours conserved ✓ • Accept: no neutrino [and no γ] emission 	2
		(c) (i) x is an electron (1) y is an antineutrino (1) clear logical reasoning based on the laws of conservation of charge and of lepton number (1)	3
	(ii) Weak	1	
Question 7 Total			[10]