



MS4
£4.00

GCE MARKING SCHEME

CHEMISTRY (NEW)
AS/Advanced

SUMMER 2009


CH1

Section A

1. (a) (i) Atomic number is the number of protons in the nucleus / in an element (e.g. 19 for potassium) [1]

(ii) Isotopes of elements have the same number of protons but different number of neutrons (e.g. chlorine has two isotopes ^{35}Cl and ^{37}Cl) / same atomic number but different mass number [1]

(b)




2. (a) (i) Measure (the volume of) hydrogen produced (using a gas syringe) / (mass of) hydrogen lost at constant time intervals [1]

(ii) Crush it into a powder / increase its surface area / heat it / stir it [1]

(b) 2 g [1]

3. 3 g / A [1]

4. (a)



(b) $\Delta H = (4 \times 412) + 612 + 436 - ((6 \times 412) + 348)$ [1]
 $= -124 \text{ kJ mol}^{-1}$ [1]

Total [10]

Section B

5. (a) (i) Correct plotting of 6 points (Allow $\pm \frac{1}{2}$ square) [3]
- (ii) In He less shielding of outer electron (1)
outweighs smaller nuclear charge (1) /
He has greater effective nuclear charge (1) /
He outer electron closer to nucleus (1)
- (Accept any two points) [2]
- (iii) Ne has greater nuclear charge /
greater number of protons (in same orbital) [1]
- (iv) N only has unpaired 2p electrons, O has two unpaired
and two paired 2p electrons / N $1s^2 2s^2 2p^3$, O $1s^2 2s^2 2p^4$ (1),
repulsion between the paired electrons makes it easier to
remove one of the electrons / takes more energy to remove
unpaired electron (1) [2]

- (b) (i)
- | Pb | C | O |
|-------------------------------|-------------------|-------------------|
| $\frac{77.5}{207}$ | $\frac{4.50}{12}$ | $\frac{18.0}{16}$ |
| 0.374 | 0.375 | 1.125 (1) |
| 1 | 1 | 3 |
| Formula = PbCO_3 (1) | | |
- [2]

- (ii) I $M_r \text{Pb}_3\text{O}_4 = (3 \times 207) + (4 \times 16) = 685$ [1]
- II Moles PbO = $\frac{134}{223} = 0.601$ (1)
- Moles $\text{Pb}_3\text{O}_4 = 0.200$ (1)
- Mass $\text{Pb}_3\text{O}_4 = 137 \text{ g}$ (1) [3]

or alternative

$$1338 \text{ g PbO gives } 1370 \text{ g Pb}_3\text{O}_4 \quad (1)$$

$$1 \text{ g PbO gives } \frac{1370}{1388} \text{ g Pb}_3\text{O}_4 \quad (1)$$

$$134 \text{ g PbO gives } 137(.2) \text{ g Pb}_3\text{O}_4 \quad (1)$$

Total [14]

6. (a) (i) It provides a new route (1)
of lower activation energy (1) [2]
- (ii) Heterogenous [1]
- (iii) I Lower temperatures could be used (1)
(which would mean) increased yield (1) /
less energy consumption (1) / lower pressure used (1) /
equilibrium could be reached faster (1)
(Accept any two points) [2]
- II More ammonia formed / equilibrium moves to right (1)
since more (gas) molecules on l.h.s. (1)
(Increases rate of reaction 1 mark) [2]
- III Equilibrium moves to right / more ammonia formed (1)
since removing ammonia decreases its concentration in
the mixture (1)
(Stops ammonia from returning to nitrogen and
hydrogen 1 mark) [2]
- (iv) Near a port / on the coast for exporting products (1),
good transport links for product (1), nearby workforce (1)


(Two valid reasons without one qualification 1 mark only) [2]
- (b) (i) $2\text{NH}_3 + \text{H}_2\text{SO}_4 \longrightarrow (\text{NH}_4)_2\text{SO}_4$ [1]
- (ii) Ammonia accepts a proton (from the acid) / ammonia has a
lone pair of electrons / ammonia neutralises the acid [1]
- (iii) % N = $28/132 \times 100$ (1)
= 21.2% (1) [2]

Total [15]

7. (a) (i) Only changes between energy levels allowed /
 electron falls from higher energy levels to lower energy levels
 (1)

Energy emitted related to frequency / $E = hf$ / the difference between any two energy levels are fixed / energy levels are quantised (1) [2]

(ii)



- Labelling of any 3 horizontal lines (1)
 Transitions going to n = 2 (1)
 Red line from n = 3 to n = 2 (1)
 (If all lines go to n = 1, accept red line from n = 2 to n = 1) [3]

(iii) Transition from n = 1 to n = ∞ [1]

$$\begin{aligned}
 (b) \quad (i) \quad A_r H &= \frac{(1 \times 99.2) + (2 \times 0.8)}{100} && (1) \\
 &= 1.008 && (1) \quad [2]
 \end{aligned}$$

(ii) Some of the hydrogen molecules are split into atoms [1]

(c) (i) Electron gun / source of electrons / heated filament [1]

(ii) Electric field / charged plates / accelerator / collimator [1]

(iii) To ensure a vacuum /
prevents collisions between sample and air molecules [1]

(d)

<i>Type</i>	<i>Nature</i>	<i>Effect on atomic number</i>
α particle	Cluster of 2 protons and 2 neutrons (1) / ${}^4_2\text{He}$ <u>nucleus</u>	Decrease by 2 (1)
β particle	Electron (1)	Increase by 1 (1)
γ radiation	Electromagnetic radiation of high energy	No effect

(Accept 'decrease' and 'increase' in 'atomic number' for 1 mark only)
[4]

Total [16]


8. (a) (i) Increases CO₂ levels / causes global warming (1)
Gas is a non renewable energy source / will run out (1) [2]

(QWC) *The information is organised clearly and coherently, using specialist vocabulary where appropriate* [1]

- (ii) Wind / hydro / biomass / solar / geothermal (1)

Rotation of blades turns turbine / falling water turns turbine / combustion steam turns turbine / sunlight on photovoltaic cell produces electricity (1)

(Accept answers in terms of energy changes) [2]



(ii) $\Delta H = (2 \times -394) + (3 \times -286) - (-278)$ (1)

$\Delta H = -1368 \text{ kJ mol}^{-1}$ (1) [2]

(iii) Energy for ethanol = $\frac{1368}{46} = 29.7 \text{ kJ g}^{-1}$ (1)

Energy for octane = $\frac{5512}{114} = 48.4 \text{ kJ g}^{-1}$ (1) [2]

- (iv) Ethanol is a renewable fuel (if obtained by fermentation) / ethanol is cheaper in countries with plentiful sugar cane growth / ethanol is more carbon neutral / ethanol burns more cleanly [1]

Total [11]

9. (a) Volumetric / graduated / standard flask [1]
- (b) 23.10 23.95 23.20 23.15 [1]
- (c) Anomalous result = 23.95 cm³
Mean = 23.15 cm³ [1]
- (d) (i) Moles HCl = $\frac{0.1 \times 23.15}{1000} = 2.315 \times 10^{-3}$ [1]
- (ii) Moles Na₂CO₃ = 1.158 × 10⁻³ [1]
- (iii) Moles in original solution = 1.158 × 10⁻² [1]
- (iv) Mass Na₂CO₃ = 1.227 g [1]
- (v) % Na₂CO₃ = 59.9 % [1]
- (Consequential marking applies)
- (e) e.g. funnel left in burette (1) / air in pipette (1) /
not reading meniscus (1) / solution in flask not mixed thoroughly (1)
/all of solid not used to make solution (1)
(Maximum 2 marks for sources of error)
If end-point overshoot, too much acid would have been added (1),
so moles (mass) carbonate calculated would have been more than
actual moles (mass) present (1) [4]
- (QWC) *Legibility of text; accuracy of spelling, punctuation and grammar,*
clarity of meaning (1)
Selection of a form and style of writing appropriate to purpose and to
complexity of subject matter (1) [2]

Total [14]

Section B Total [70]