

2007 FP2 Adapted


1. Obtain the general solution of the differential equation

$$x \frac{dy}{dx} + 2y = \cos x, \quad x > 0,$$

giving your answer in the form $y = f(x)$.

(Total 8 marks)

- 2.



The diagram above shows a sketch of the curve with equation

$$y = \frac{x^2 - 1}{|x + 2|}, \quad x \neq -2.$$

The curve crosses the x -axis at $x = 1$ and $x = -1$ and the line $x = -2$ is an asymptote of the curve.

- (a) Use algebra to solve the equation $\frac{x^2 - 1}{|x + 2|} = 3(1 - x)$.

(6)

(b) Hence, or otherwise, find the set of values of x for which

$$\frac{x^2 - 1}{|x + 2|} < 3(1 - x).$$

(3)
(Total 9 marks)

3. A scientist is modelling the amount of a chemical in the human bloodstream. The amount x of the chemical, measured in mg l^{-1} , at time t hours satisfies the differential equation

$$2x \frac{d^2x}{dt^2} - 6 \left(\frac{dx}{dt} \right)^2 = x^2 - 3x^4, \quad x > 0.$$

(a) Show that the substitution $y = \frac{1}{x^2}$ transforms this differential equation into

$$\frac{d^2y}{dt^2} + y = 3. \quad \boxed{I}$$

(5)

(b) Find the general solution of differential equation \boxed{I} .

(4)

Given that at time $t = 0$, $x = \frac{1}{2}$ and $\frac{dx}{dt} = 0$,


(c) find an expression for x in terms of t ,

(4)

(d) write down the maximum value of x as t varies.

(1)
(Total 14 marks)

4.



The diagram above shows a sketch of the curve C with polar equation

$$r = 4\sin\theta\cos^2\theta, \quad 0 \leq \theta < \frac{\pi}{2}.$$

The tangent to C at the point P is perpendicular to the initial line.

- (a) Show that P has polar coordinates $\left(\frac{3}{2}, \frac{\pi}{6}\right)$.

(6)

The point Q on C has polar coordinates $\left(\sqrt{2}, \frac{\pi}{4}\right)$.

The shaded region R is bounded by OP , OQ and C , as shown in the diagram above.

- (b) Show that the area of R is given by

$$\int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \left(\sin^2 2\theta \cos 2\theta + \frac{1}{2} - \frac{1}{2} \cos 4\theta \right) d\theta$$

(3)

- (c) Hence, or otherwise, find the area of R , giving your answer in the form $a + b\pi$, where a and b are rational numbers.

(5)

(Total 14 marks)

5. Find the set of values of x for which

$$\frac{x+1}{2x-3} < \frac{1}{x-3}$$

(Total 7 marks)

6. $\frac{dy}{dx} - y \tan x = 2 \sec^3 x.$

Given that $y = 3$ at $x = 0$, find y in terms of x

(Total 7 marks)

7. For the differential equation

$$\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = 2x(x+3),$$

find the solution for which at $x = 0$, $\frac{dy}{dx} = 1$ and $y = 1$.

(Total 12 marks)

8. (a) Sketch the curve C with polar equation

$$r = 5 + \sqrt{3} \cos \theta, \quad 0 \leq \theta \leq 2\pi. \quad (2)$$

- (b) Find the polar coordinates of the points where the tangents to C are parallel to the initial line $\theta = 0$. Give your answers to 3 significant figures where appropriate.

(6)

- (c) Using integration, find the area enclosed by the curve C , giving your answer in terms of π .

(6)

(Total 14 marks)

9.

$$\frac{dy}{dx} = ye^{x^2}.$$

It is given that $y = 0.2$ at $x = 0$.

- (a) Use the approximation $\frac{y_1 - y_0}{h} \approx \left(\frac{dy}{dx}\right)_0$, with $h = 0.1$, to obtain an estimate of the value of y at $x = 0.1$.

(2)

- (b) Use your answer to part (a) and the approximation $\frac{y_2 - y_0}{2h} \approx \left(\frac{dy}{dx}\right)_1$, with $h = 0.1$, to obtain an estimate of the value of y at $x = 0.2$.

Gives your answer to 4 decimal places.

(3)

(Total 5 marks)

10.

$$(1 - x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + 2y = 0.$$

At $x = 0$, $y = 2$ and $\frac{dy}{dx} = -1$.

- (a) Find the value of $\frac{d^3y}{dx^3}$ at $x = 0$.

(3)

- (b) Express y as a series in ascending powers of x , up to and including the term in x^3 .

(4)

(Total 7 marks)

11. (a) Given that $z = \cos \theta + i \sin \theta$, use de Moivre's theorem to show that

$$z^n + \frac{1}{z^n} = 2 \cos n\theta. \quad (2)$$

- (b) Express $32\cos^6\theta$ in the form $p\cos 6\theta + q\cos 4\theta + r\cos 2\theta + s$, where p, q, r and s are integers. (5)

- (c) Hence find the exact value of

$$\int_0^{\frac{\pi}{3}} \cos^6 \theta \, d\theta. \quad (4)$$

(Total 11 marks)

12. The transformation T from the z -plane, where $z = x + iy$, to the w -plane, where $w = u + iv$, is given by

$$w = \frac{z+i}{z}, \quad z \neq 0.$$

- (a) The transformation T maps the points on the line with equation $y = x$ in the z -plane, other than $(0, 0)$, to points on a line l in the w -plane. Find a cartesian equation of l . (5)

- (b) Show that the image, under T , of the line with equation $x + y + 1 = 0$ in the z -plane is a circle C in the w -plane, where C has cartesian equation

$$u^2 + v^2 - u + v = 0. \quad (7)$$

- (c) On the same Argand diagram, sketch l and C . (3)
- (Total 15 marks)**