

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	



General Certificate of Education  
Advanced Level Examination  
January 2011

# Mathematics

# MPC4

## Unit Pure Core 4

Monday 24 January 2011 9.00 am to 10.30 am

**For this paper you must have:**

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

### Time allowed

- 1 hour 30 minutes

### Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer the questions in the spaces provided. Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

### Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.



J A N 1 1 M P C 4 0 1

Answer **all** questions in the spaces provided.

**1 (a)** Express  $2 \sin x + 5 \cos x$  in the form  $R \sin(x + \alpha)$ , where  $R > 0$  and  $0^\circ < \alpha < 90^\circ$ .  
Give your value of  $\alpha$  to the nearest  $0.1^\circ$ . *(3 marks)*

**(b) (i)** Write down the maximum value of  $2 \sin x + 5 \cos x$ . *(1 mark)*

**(ii)** Find the value of  $x$  in the interval  $0^\circ \leq x \leq 360^\circ$  at which this maximum occurs,  
giving the value of  $x$  to the nearest  $0.1^\circ$ . *(2 marks)*

QUESTION  
PART  
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- 2 (a)** The polynomial  $f(x)$  is defined by  $f(x) = 9x^3 + 18x^2 - x - 2$ .
- (i) Use the Factor Theorem to show that  $3x + 1$  is a factor of  $f(x)$ . (2 marks)
  - (ii) Express  $f(x)$  as a product of three linear factors. (3 marks)
  - (iii) Simplify  $\frac{9x^3 + 21x^2 + 6x}{f(x)}$ . (3 marks)
- (b)** When the polynomial  $9x^3 + px^2 - x - 2$  is divided by  $3x - 2$ , the remainder is  $-4$ .  
Find the value of the constant  $p$ . (2 marks)

QUESTION PART REFERENCE	
-------------------------------	--



**3 (a)** Express  $\frac{3 + 9x}{(1 + x)(3 + 5x)}$  in the form  $\frac{A}{1 + x} + \frac{B}{3 + 5x}$ , where  $A$  and  $B$  are integers. (3 marks)

**(b)** Hence, or otherwise, find the binomial expansion of  $\frac{3 + 9x}{(1 + x)(3 + 5x)}$  up to and including the term in  $x^2$ . (7 marks)

**(c)** Find the range of values of  $x$  for which the binomial expansion of  $\frac{3 + 9x}{(1 + x)(3 + 5x)}$  is valid. (2 marks)

QUESTION  
PART  
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



4 A curve is defined by the parametric equations

$$x = 3e^t, \quad y = e^{2t} - e^{-2t}$$

(a) (i) Find the gradient of the curve at the point where  $t = 0$ . (3 marks)

(ii) Find an equation of the tangent to the curve at the point where  $t = 0$ . (1 mark)

(b) Show that the cartesian equation of the curve can be written in the form

$$y = \frac{x^2}{k} - \frac{k}{x^2}$$

where  $k$  is an integer. (2 marks)

QUESTION  
PART  
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



5 A model for the radioactive decay of a form of iodine is given by

$$m = m_0 2^{-\frac{1}{8}t}$$

The mass of the iodine after  $t$  days is  $m$  grams. Its initial mass is  $m_0$  grams.

(a) Use the given model to find the mass that remains after 10 grams of this form of iodine have decayed for 14 days, giving your answer to the nearest gram. (2 marks)

(b) A mass of  $m_0$  grams of this form of iodine decays to  $\frac{m_0}{16}$  grams in  $d$  days.

Find the value of  $d$ . (2 marks)

(c) After  $n$  days, a mass of this form of iodine has decayed to less than 1% of its initial mass.

Find the minimum integer value of  $n$ . (3 marks)

QUESTION  
PART  
REFERENCE

A large rectangular area containing horizontal dotted lines, intended for the student to provide answers and working for parts (a), (b), and (c).



**6 (a) (i)** Given that  $\tan 2x + \tan x = 0$ , show that  $\tan x = 0$  or  $\tan^2 x = 3$ . *(3 marks)*

**(ii)** Hence find all solutions of  $\tan 2x + \tan x = 0$  in the interval  $0^\circ < x < 180^\circ$ . *(1 mark)*

**(b) (i)** Given that  $\cos x \neq 0$ , show that the equation

$$\sin 2x = \cos x \cos 2x$$

can be written in the form

$$2 \sin^2 x + 2 \sin x - 1 = 0 \quad \text{\textit{(3 marks)}}$$

**(ii)** Show that all solutions of the equation  $2 \sin^2 x + 2 \sin x - 1 = 0$  are given by

$$\sin x = \frac{\sqrt{3} - 1}{p}, \text{ where } p \text{ is an integer.} \quad \text{\textit{(3 marks)}}$$

QUESTION  
PART  
REFERENCE



**7 (a) (i)** Solve the differential equation  $\frac{dx}{dt} = \sqrt{x} \sin\left(\frac{t}{2}\right)$  to find  $x$  in terms of  $t$ . (3 marks)

**(ii)** Given that  $x = 1$  when  $t = 0$ , show that the solution can be written as

$$x = (a - \cos bt)^2$$

where  $a$  and  $b$  are constants to be found. (3 marks)

**(b)** The height,  $x$  metres, above the ground of a car in a fairground ride at time  $t$  seconds is modelled by the differential equation  $\frac{dx}{dt} = \sqrt{x} \sin\left(\frac{t}{2}\right)$ .

The car is 1 metre above the ground when  $t = 0$ .

**(i)** Find the greatest height above the ground reached by the car during the ride. (2 marks)

**(ii)** Find the value of  $t$  when the car is first 5 metres above the ground, giving your answer to one decimal place. (2 marks)

QUESTION  
PART  
REFERENCE

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





**8** The coordinates of the points  $A$  and  $B$  are  $(3, -2, 4)$  and  $(6, 0, 3)$  respectively.

The line  $l_1$  has equation  $\mathbf{r} = \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix} + \lambda \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}$ .


**(a) (i)** Find the vector  $\overrightarrow{AB}$ . (2 marks)

**(ii)** Calculate the acute angle between  $\overrightarrow{AB}$  and the line  $l_1$ , giving your answer to the nearest  $0.1^\circ$ . (4 marks)

**(b)** The point  $D$  lies on  $l_1$  where  $\lambda = 2$ . The line  $l_2$  passes through  $D$  and is parallel to  $AB$ .

**(i)** Find a vector equation of line  $l_2$  with parameter  $\mu$ . (2 marks)

**(ii)** The diagram shows a symmetrical trapezium  $ABCD$ , with angle  $DAB$  equal to angle  $ABC$ .



The point  $C$  lies on line  $l_2$ . The length of  $AD$  is equal to the length of  $BC$ .

Find the coordinates of  $C$ . (6 marks)

QUESTION  
PART  
REFERENCE

.....

.....

.....

.....

.....

.....

.....

