

Version1.0



**General Certificate of Education (A-level)
January 2011**

Statistics

SS03

(Specification 6380)

Statistics 3

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2011 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
√ or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
-x EE	deduct x marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

Q	Solution						Marks	Total	Comments
1(a)	Male	A	B	C	D	E	M1	6	attempt at ranks 18 correct (can be reversed) all correct — can be reversed alternative $d = 10, 7, 4, 8, 5, 2, 1, 2\frac{1}{2}, 3\frac{1}{2}, 7, 10, 10$ $\sum d^2 = 526\frac{1}{2}$ B1 $r_s = 1 - \frac{6 \times 526.5}{12 \times 143} = -0.841$ M1, A1 SC4 -0.84 (no method seen) SC2 -0.8 (no method seen) SC4 0.846 (no ties) SC4 +0.844 (inconsistent ranks)
	x rank	1	2	3	4	5	M1		
	y rank	11	9	7	12	10			
	Male	F	G	H	I	J	A1		
	x rank	6	7	8½	8½	10			
	y rank	4	8	6	5	3			
	Male	K	L						
	x rank	11	12						
	y rank	1	2						
		$r_s = -0.844$ (3 sf from calc)							
(b)	H_0 Rank orders number of average number of hours slept and diastolic blood pressure are independent. H_1 Rank orders are not independent. 2 tail 1% $cv = \pm 0.7273$ test stat $r_s = -0.844$ (or -0.841) $r_s < -0.7273$ or $ r_s > cv $ Reject H_0 . Significant evidence at 1% level to suggest an association between rank orders number of average number of hours slept and diastolic blood pressure. Results suggests that adult males who sleep less on average tend to have a higher diastolic blood pressure.						B1		
							B1	for cv — ignore sign	
							M1	for comparison ts/cv Must be consistent = or –	
							E1	4 in context	
Total							10		

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments																																														
2(a)	$H_0 \eta_{\text{difference}} = 0$ $H_1 \eta_{\text{difference}} < 0$ 1 tail 5%	B1		Allow μ ; condone no difference' mentioned or $\eta_{\text{difference}} > 0$ if consistent																																														
	<table border="1"> <thead> <tr> <th rowspan="2">Burglar pair</th> <th rowspan="2">Difference 1994 –2004</th> <th colspan="2">Rank</th> </tr> <tr> <th>–</th> <th>+</th> </tr> </thead> <tbody> <tr><td>A</td><td>–4</td><td>2</td><td></td></tr> <tr><td>B</td><td>–8</td><td>7</td><td></td></tr> <tr><td>C</td><td>+1</td><td></td><td>1</td></tr> <tr><td>D</td><td>–5</td><td>3</td><td></td></tr> <tr><td>E</td><td>–9</td><td>8</td><td></td></tr> <tr><td>F</td><td>+7</td><td></td><td>5½</td></tr> <tr><td>G</td><td>–10</td><td>9</td><td></td></tr> <tr><td>H</td><td>–7</td><td>5½</td><td></td></tr> <tr><td>I</td><td>–6</td><td>4</td><td></td></tr> <tr><td>J</td><td>–11</td><td>10</td><td></td></tr> </tbody> </table>	Burglar pair	Difference 1994 –2004	Rank		–	+	A	–4	2		B	–8	7		C	+1		1	D	–5	3		E	–9	8		F	+7		5½	G	–10	9		H	–7	5½		I	–6	4		J	–11	10		M1		for differences
	Burglar pair			Difference 1994 –2004	Rank																																													
		–	+																																															
	A	–4	2																																															
	B	–8	7																																															
	C	+1		1																																														
	D	–5	3																																															
	E	–9	8																																															
	F	+7		5½																																														
	G	–10	9																																															
	H	–7	5½																																															
	I	–6	4																																															
J	–11	10																																																
		m1		for ranks																																														
	Rank totals $T_- = 48\frac{1}{2}$ $T_+ = 6\frac{1}{2}$ Test stat $T = 6\frac{1}{2}$	m1 A1		For totals (dep ranks) One total correct																																														
	$n = 10$ critical value = 11	B1		For cv																																														
	$T < cv$	M1																																																
	Reject H_0 . There is significant evidence to suggest that sentence lengths have increased since 1994	A1 E1	9	In context																																														
(b)	Matched pairs design will eliminate individual differences between types of burglar/types of burglary and will therefore reduce experimental error and make the test more likely to detect a difference if one exists.	E1		In context																																														
		B1	2	General terms																																														
			11																																															

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments
3(a)(i)	H ₀ Type of accident is independent of whether HGV British registered or foreign registered H ₁ Type of accident is not independent of whether HGV British registered or foreign registered 1 tail 1%	B1		
		M1		E method for 3 correct
		A1		For all E correct to 1 dp
				[Alt $\frac{(O-E)^2}{E}$ clearly > 9.21 at early stage]
		m1		ts sum with correct denominators
		A1		For ts in range 260 ~ 270
		B1		For cv
		m1		For comparison ts/cv
		A1		
		E1	9	[Alt Allow clear explanation ref 266 being so large therefore significant as alternative]
(ii)	Sideswipe accidents involving changing lanes to left far less likely than expected for foreign registered HGVs and accidents involving changing lanes to right far more likely than expected for foreign registered HGVs.	E1		Or E1 relevant comment on overtaking.
		E1	2	Must identify association sideswipe/changing right with foreign HGV for E1, E1
(b)(i)		B1		Labels correct
		M1		2 correct
		A1	3	

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments									
<p>3(b)(ii)</p>	<p>H_0 Age of driver is independent of whether prosecution resulted. H_1 Age of driver is not independent of whether prosecution resulted. 1 tail 5%</p> <table border="1" data-bbox="240 405 692 629"> <tr> <td>Exp values</td> <td>Prosecution resulted</td> <td>No prosecution</td> </tr> <tr> <td>35 years or under</td> <td>10.36</td> <td>17.64</td> </tr> <tr> <td>Over 35 years</td> <td>26.64</td> <td>45.36</td> </tr> </table> $ts = \sum \frac{(O - E - 0.5)^2}{E} =$ $\frac{1.86^2}{10.36} + \frac{1.86^2}{17.64} + \frac{1.86^2}{26.64} + \frac{1.86^2}{45.36} =$ <p style="text-align: right;">0.736</p> <p>cv df = 1 5% cv = 3.841 ts < 3.841</p> <p>Accept H_0 No significant evidence to suggest an association. Conclude that age of driver is independent of whether prosecution resulted</p>	Exp values	Prosecution resulted	No prosecution	35 years or under	10.36	17.64	Over 35 years	26.64	45.36	<p>B1</p> <p>M1</p> <p>M1 m1</p> <p>A1</p> <p>B1</p> <p>E1</p>	<p>7</p>	<p>For E values method</p> <p>For ts for Yates' corr</p> <p>For ts 0.70 ~ 0.77</p> <p>For cv</p>
Exp values	Prosecution resulted	No prosecution											
35 years or under	10.36	17.64											
Over 35 years	26.64	45.36											
	Total		21										

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments
4	<p>H_0 samples from identical populations</p> <p>H_1 samples not from identical populations: taste better on average for pods produced using new method</p> <p>1 tail 5%</p> <p>Current method ranks 10 12 6 8 11 5</p> <p>New method ranks 3 2 7 1 4 9</p> <p>$T_{\text{current}} = 52$ $T_{\text{new}} = 26$</p> <p>$U_{\text{current}} = 52 - \frac{(6 \times 7)}{2} = 31$</p> <p>$U_{\text{new}} = 26 - \frac{(6 \times 7)}{2} = 5$</p> <p>test stat = 5 (lower)</p> <p>$n = 6, m = 6$ cv = lower tail 7</p> <p>Since $5 < 7$, reject H_0</p> <p>Significant evidence to suggest that populations are not identical and that the taste is better, on average, for pods produced using new method.</p>	<p>B1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>E1</p>	<p>9</p>	<p>Sorting into 2 groups</p> <p>Totals</p> <p>Method for U</p> <p>Either U correct</p> <p>cv</p> <p>Comparison correct cv and ts (can be upper tail)</p>
	Total		9	

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments	
5(a)	H ₀ Samples are taken from identical populations H ₁ Samples are not taken from identical populations 1% sig level	B1		or H ₀ $\eta_A = \eta_B = \eta_C = \eta_D = \eta_E$ H ₁ at least two of $\eta_A, \eta_B, \eta_C, \eta_D, \eta_E$ do differ	
	$T_A = 75$ $T_B = 99$ $T_C = 26$ $T_D = 31$ $T_E = 94$ $n_A = 5$ $n_B = 5$ $n_C = 5$ $n_D = 5$ $n_E = 5$	M1 A1		totals any one correct	
	$\sum_{i=1}^m \frac{T_i^2}{n_i} = \frac{75^2}{5} + \frac{99^2}{5} + \frac{26^2}{5} + \frac{31^2}{5} + \frac{94^2}{5}$ $= 5179.8$	m1		$\sum_{i=1}^m \frac{T_i^2}{n_i}$	
	$H = \frac{12}{25 \times 26} \times 5179.8 - (3 \times 26) = 17.63$	m1		test stat: $H = \frac{12}{N(N+1)} \sum_{i=1}^m \frac{T_i^2}{n_i} - 3(N+1)$ 17.0 ~ 18.4	
	Critical value from $\chi_4^2 = 13.277$ $H > 13.277$	A1 B1 M1		for cv for comparison	
	Sig evidence to reject H ₀ and conclude that samples are not from identical populations. At least 2 average acidity levels are different.	E1	9		
	(b) Variety B has highest total of ranks so if a low acidity beer is desirable, this variety would be the best choice.	B1 E1	2	Identification of B Explained	
	(c) Conclusion only shows that Variety B differs significantly from Variety C (highest and lowest). However, Variety B and Variety E have similar acidity level ranks. Thus Variety E is a sensible choice if popular with customers.	E1 E1	2		
	Total			13	

Mark Scheme – General Certificate of Education (A-level) Statistics – Statistics 3 – January 2011

SS03(cont)

Q	Solution	Marks	Total	Comments
6(a)(i)	H ₀ Women like the taste of both recipes equally, on average	B1	2	1 tail correct
	H ₁ On average, women prefer the taste of the new recipe.	B1		context/wording correct — mention women
(ii)	1 tail test 5% level			H ₀ no preference H ₁ preference B1 only
	test stat 10+ or 5– B(15, 0.5) model	B1 M1		for test stat for model B(15, 0.5) seen
	P (≥ 10+) = P (≤ 5–) = 0.151 Since 0.151 > 0.05 for 1 tail test	M1		correct probability and comparison with 0.05
(b)	Accept H ₀ No sig evidence to suggest adult females prefer new recipe	A1 E1	5	In context
	B(30, 0.5) model 1 tail 5% level	M1		Use of B(30. 0.5) method must be seen
	P (≥ n +) < 0.05 required from tables,	M1		Comparison of B(30, 0.5) probability with 0.05
	P(≤ 10 –) = P(≥ 20+) = 0.0494 P(≤ 11 –) = P(≥ 19+) = 0.1002	M1		Correct B(30, 0.5) probability seen
	minimum number therefore 20 adult females out of the 30 to prefer the new recipe.	A1	4	Or equivalent
	Total		11	
	TOTAL		75	